

Publications - First Author

1. **Iida O**, Takahara M. What does Intravascular Ultrasound Illustrate? *J Atheroscler Thromb*. 2016 Nov 29. [Epub ahead of print]
2. **Iida O**, Takahara M, Soga Y, Hirano K, Yamauchi Y, Zen K, Yokoi H, Uematsu M; ZEPHYR investigators. Incidence and its characteristics of repetition of reintervention after drug-eluting stent implantation for femoropopliteal lesion. *J Vasc Surg*. 2016;64:1691-1695.e1.
3. **Iida O**, Takahara M, Soga Y, Hirano K, Yamauchi Y, Zen K, Kawasaki D, Nanto S, Yokoi H, Uematsu M; ZEPHYR Investigators. The Characteristics of In-Stent Restenosis After Drug-Eluting Stent Implantation in Femoropopliteal Lesions and 1-Year Prognosis After Repeat Endovascular Therapy for These Lesions. *JACC Cardiovasc Interv*. 2016;9:828-34.
4. **Iida O**, Nakamura M. Reply: 3-Year Outcomes of the OLIVE Registry, a Prospective Multicenter Study of Patients with Critical Limb Ischemia. *JACC Cardiovasc Interv*. 2016;9:202-3.
5. **Iida O**, Nakamura M, Yamauchi Y, Fukunaga M, Yokoi Y, Yokoi H, Soga Y, Zen K, Suematsu N, Inoue N, Suzuki K, Hirano K, Shintani Y, Miyashita Y, Urasawa K, Kitano I, Tsuchiya T, Kawamoto K, Yamaoka T, Uesugi M, Shinke T, Oba Y, Ohura N, Uematsu M, Takahara M, Hamasaki T, Nanto S; OLIVE Investigators. 3-Year Outcomes of the OLIVE Registry, a Prospective Multicenter Study of Patients With Critical Limb Ischemia: A Prospective, Multi-Center, Three-Year Follow-Up Study on Endovascular Treatment for Infra-Inguinal Vessel in Patients With Critical Limb Ischemia. *JACC Cardiovasc Interv*. 2015;8:1493-502.
6. **Iida O**, Takahara M, Soga Y, Nakano M, Yamauchi Y, Zen K, Kawasaki D, Nanto S, Yokoi H, Uematsu M; ZEPHYR Investigators. 1-Year Results of the ZEPHYR (Zilver PTX for the Femoral Artery and Proximal Popliteal Artery) Registry: Predictors of Restenosis. *JACC Cardiovasc Interv*. 2015;8:1105-1112.
7. **Iida O**, Takahara M, Soga Y, Yamauchi Y, Hirano K, Tazaki J, Yamaoka T, Suematsu N, Suzuki K, Shintani Y, Miyashita Y, Uematsu M. Impact of angiosome-oriented revascularization on clinical outcomes in critical limb ischemia patients without concurrent wound infection and diabetes. *J Endovasc Ther*. 2014;21:607-15.
8. **Iida O**, Takahara M, Soga Y, Suzuki K, Hirano K, Kawasaki D, Shintani Y, Suematsu N, Yamaoka T, Nanto S, Uematsu M. Efficacy of Intravascular Ultrasound in Femoropopliteal Stenting for Peripheral Artery Disease With TASC II Class A to C Lesions. *J Endovasc Ther*. 2014;21:485-492.
9. **Iida O**, Takahara M, Soga Y, Suzuki K, Hirano K, Kawasaki D, Shintani Y, Suematsu N, Yamaoka T, Nanto S, Uematsu M. Shared and Differential Factors Influencing Restenosis Following Endovascular Therapy Between TASC (Trans-Atlantic Inter-Society Consensus) II Class A to C and D Lesions in the Femoropopliteal Artery. *JACC Cardiovasc Interv*. 2014;7:792-8.

10. **Iida O**, Soga Y, Takahara M, Kawasaki D, Yamauchi Y, Suzuki K, Hirano K, Hirano K, Kamoi D, Tazaki J, Higashitani M, Shintani Y, Yamaoka T, Okazaki S, Suematsu N, Tsuchiya T, Miyashita Y, Shinozaki N, Takahashi H, Uematsu M. Perioperative Complications After Aortoiliac Stenting: Associated Factors and Impact on Follow-up Cardiovascular Prognosis. *Eur J Vasc Endovasc Surg.* 2014;47:131-138.
11. **Iida O**, Takahara M, Soga Y, Yamauchi Y, Hirano K, Tazaki J, Yamaoka T, Suematsu N, Suzuki K, Shintani Y, Miyashita Y, Uematsu M. Worse limb prognosis for indirect vs. direct endovascular revascularization only in patients with critical limb ischemia complicated with wound infection and diabetes mellitus. *Eur J Vasc Endovasc Surg.* 2013;46:575-82.
12. **Iida O**, Soga Y, Takahara M, Kawasaki D, Yamauchi Y, Suzuki K, Hirano K, Ryoji K, Kamoi D, Tazaki J, Higashitani M, Shintani Y, Yamaoka T, Okazaki S, Suematsu N, Tsuchiya T, Miyashita Y, Shinozaki N, Takahashi H, Uematsu M. Efficacy of the S.M.A.R.T. Control vs. other stents for aortoiliac occlusive disease in contemporary clinical practice. *J Endovasc Ther.* 2013;20:431-9.
13. **Iida O**, Yokoi H, Soga Y, Inoue N, Suzuki K, Yokoi Y, Kawasaki D, Zen K, Urasawa K, Shintani Y, Miyamoto A, Hirano K, Miyashita Y, Tsuchiya T, Shinozaki N, Nakamura M, Isshiki T, Hamasaki T, Nanto S; on behalf of the STOP-IC investigators. Cilostazol Reduces Angiographic Restenosis After Endovascular Therapy for Femoropopliteal Lesions in the Sufficient Treatment of Peripheral Intervention by Cilostazol Study. *Circulation.* 2013;127:2307-15.
14. **Iida O**, Soga Y, Yamauchi Y, Hirano K, Kawasaki D, Yamaoka T, Takahara M, Uematsu M. Clinical efficacy of endovascular therapy for patients with critical limb ischemia attributable to pure isolated infrapopliteal lesions. *J Vasc Surg.* 2013;57:974-981.
15. **Iida O**, Nakamura M, Miyamoto A, Kawasaki D, Yokoi Y, Yokoi H, Zen K, Hirano K, Suematsu N, Inoue N, Shintani Y, Miyashita Y, Urasawa K, Kitano I, Yamaoka T, Murakami T, Uesugi M, Tsuchiya T, Shinke T, Oba Y, Ohura N, Hamasaki T, Nanto S on behalf of the OLIVE investigators. Endovascular Treatment for Infringuinal Vessels in Patients with Critical Limb Ischemia: OLIVE Registry, a Prospective, Multicenter Study in Japan with 12-month Follow-up. *Circ Cardiovasc Interv.* 2013;6:68-76.
16. **Iida O**, Soga Y, Kawasaki D, Hirano K, Yamaoka T, Suzuki K, Miyashita Y, Yokoi H, Takahara M, Uematsu M. Angiographic restenosis and its clinical impact after infrapopliteal angioplasty. *Eur J Vasc Endovasc Surg.* 2012;44:425-31.
17. **Iida O**, Soga Y, Hirano K, Kawasaki D, Suzuki K, Miyashita Y, Nanto S, Uematsu M. Anatomical Predictors of Major Adverse Limb Events after Infrapopliteal Angioplasty for Patients with Critical Limb Ischaemia due to Pure Isolated Infrapopliteal Lesions. *Eur J Vasc Endovasc Surg.* 2012;44:318-24.
18. **Iida O**, Soga Y, Hirano K, Kawasaki D, Suzuki K, Miyashita Y, Nanto S, Uematsu M. Midterm Outcomes and Risk Stratification after Endovascular Therapy for Patients with Critical Limb Ischaemia due to Isolated Below-the-knee Lesions. *Eur J Vasc Endovasc Surg.* 2012;43:313-21.

19. **Iida O**, Soga Y, Hirano K, Kawasaki D, Suzuki K, Miyashita Y, Terashi H, Uematsu M. Long-term results of direct and indirect endovascular revascularization based on the angiosome concept in patients with critical limb ischemia presenting with isolated below-the-knee lesions. *J Vasc Surg.* 2012;55:363-370.
20. **Iida O**, Soga Y, Hirano K, Suzuki K, Yokoi H, Nobuyoshi M, Muramatsu T, Inoue N, Nanto S, Uematsu M. Long-term outcomes and risk stratification of patency following nitinol stenting in the femoropopliteal segment: retrospective multicenter analysis. *J Endovasc Ther.* 2011;18:753–761.
21. **Iida O**, Uematsu M, Soga Y, Hirano K, Suzuki K, Yokoi H, Muramatsu T, Inoue N, Nanto S, Nagata S. Timing of the restenosis following nitinol stenting in the superficial femoral artery and the factors associated with early and late restenoses. *Catheter Cardiovasc Interv.* 2011;78:611-7.
22. **Iida O**, Soga Y, Hirano K, Okamoto S, Dohi T, Uematsu M, Yokoi H, Nobuyoshi M, Muramatsu T, Fujita M, Nanto S, Nagata S. Retrospective multicentre analysis of S.M.A.R.T. vs. Luminexx nitinol stent implantation for superficial femoral artery lesions (REAL SL) Registry. 5 years' experience. *Circ J.* 2011;75:421-7.
23. **Iida O**, Nanto S, Uematsu M, Ikeoka K, Okamoto S, Dohi T, Fujita M, Terashi H, Nagata S. Importance of the angiosome concept for endovascular therapy in patients with critical limb ischemia. *Catheter Cardiovasc Interv.* 2010;75:830-6.
24. **Iida O**, Nanto S, Uematsu M, Ikeoka K, Okamoto S, Dohi T, Fujita M, Nagata S. Long-term results of endovascular therapy with nitinol stent implantation for TASC II A/B femoro-popliteal artery lesions: 4 years' experience. *Circ J.* 2009;73:2143-7.
25. **Iida O**, Nanto S, Uematsu M, Ikeoka K, Okamoto S, Nagata S. Influence of stent fracture on the long-term patency in the femoro-popliteal artery: experience of 4 years. *JACC Cardiovasc Interv.* 2009;2:665-71.
26. **Iida O**, Nanto S, Uematsu M, Morozumi T, Ikeoka K, Terashi H, Nagata S. Novel Infra-popliteal Bi-directional Angioplasty in a Patient with Chronic Total Occlusion of the Tibial Artery Complicating Critical Lower Limb Ischemia. *Jpn J Interv Cardiol.* 2008;23:365-9.
27. **Iida O**, Nanto S, Uematsu M, Morozumi T, Kitakaze M, Nagata S. Cilostazol reduces restenosis after endovascular therapy in patients with femoropopliteal lesions. *J Vasc Surg.* 2008;48:144-9.
28. **Iida O**, Nanto S, Uematsu M, Morozumi T, Akahori H, Nagata S. Endovascular therapy for limb salvage in a case of critical lower limb ischemia resulting from fibromuscular dysplasia. *J Vasc Surg.* 2007;46:803-7.
29. **Iida O**, Nanto S, Uematsu M, Morozumi T, Kotani J, Awata M, Onishi T, Ito N, Sera F, Minamiguchi H, Akahori H, Nagata S. Effect of exercise on frequency of stent fracture in the superficial femoral artery. *Am J Cardiol.* 2006;98:272-4.

30. **Iida O**, Nanto S, Uematsu M, Morozumi T, Kotani J, Awata M, Onishi T, Ito N, Oshima F, Minamiguchi H, Kitakaze M, Nagata S. Cilostazol reduces target lesion revascularization after percutaneous transluminal angioplasty in the femoropopliteal artery. *Circ J.* 2005;69:1256-9.

Publications - Co-Author

1. Masuda M, Fujita M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Kanda T, Sunaga A, Tsujimura T, Matsuda Y, Ohashi T, Uematsu M. Comparison of Left Atrial Voltage Between Sinus Rhythm and atrial Fibrillation in Association with Electrogram Wave Form. *Pacing Clin Electrophysiol*. 2017 Feb 16. [Epub ahead of print]
2. Utsunomiya M, Takahara M, **Iida O**, Yamauchi Y, Kawasaki D, Yokoi Y, Soga Y, Ohura N, Nakamura M; OLIVE Investigators. Wound Blush Obtainment Is the Most Important Angiographic Endpoint for Wound Healing. *JACC Cardiovasc Interv*. 2017;10:188-194.
3. Zen K, Takahara M, **Iida O**, Soga Y, Kawasaki D, Nanto S, Yokoi H, Matoba S; ZEPHYR Investigators. Drug-eluting stenting for femoropopliteal lesions, followed by cilostazol treatment, reduces stent restenosis in patients with symptomatic peripheral artery disease. *J Vasc Surg*. 2017 Jan 7. [Epub ahead of print]
4. Suemitsu K, **Iida O**, Shiraki T, Suemitsu S, Murakami M, Miyamoto M, Izumi M, Nakanishi T. Predicting loss of patency after forearm loop arteriovenous graft. *J Vasc Surg*. 2016;64:395-401.
5. Shiraki T, **Iida O**, Takahara M, Soga Y, Mii S, Okazaki J, Kuma S, Yamaoka T, Kamoi D, Shintani Y, Ishikawa T, Kitano I, Uematsu M. Comparison of Clinical Outcomes after Surgical and Endovascular Revascularization in Hemodialysis Patients with Critical Limb Ischemia. *J Atheroscler Thromb*. 2016 Oct 13. [Epub ahead of print]
6. Rundback JH, Armstrong EJ, Contos B, **Iida O**, Jacobs D, Jaff MR, Matsumoto AH, Mills JL Sr, Montero-Baker M, Pena C, Tallian A, Uematsu M, Wilkins LR, Shishehbor MH. Key Concepts in Critical Limb Ischemia: Selected Proceedings from the 2015 Vascular Interventional Advances Meeting. *Ann Vasc Surg*. 2017;38:191-205.
7. Suzuki K, Takahara M, Shintani Y, Tanaka A, Soga Y, Yamaoka T, Tosaka A, Sasaki S, Kawasaki D, Tsuchiya T, Kozuki A, **Iida O**, Retrospective Multicenter Comparison of S.M.A.R.T. CONTROL and MISAGO Stents in Treatment of Femoropopliteal Lesions. *J Vasc Interv Radiol*. 2016;27:1642-1649.
8. Ishihara T, Takahara M, **Iida O**, Soga Y, Hirano K, Yamauchi Y, Zen K, Kawasaki D, Nanto S, Yokoi H, Uematsu M; ZEPHYR Investigators. Comparable 2-Year Restenosis Rates Following Subintimal and Intraluminal Drug-Eluting Stent Implantation for Femoropopliteal Chronic Total Occlusion. *J Endovasc Ther*. 2016;23:889-895.
9. Tsuchiya T, Takamura T, Soga Y, **Iida O**, Hirano K, Suzuki K, Yamaoka T, Miyashita Y, Kitayama M, Kajinami K. Clinical impact and risk stratification of balloon angioplasty for femoropopliteal disease in nitinol stenting era: Retrospective multicenter study using propensity score matching analysis. *SAGE Open Med*. 2016 Jul 18;4:2050312116660116.
10. Yanaka K, Ishihara T, **Iida O**, Okamoto S, Fujita M, Masuda M, Nanto K, Shiraki T, Kanda T, Uematsu M. Yellow Neointima Following Stent Implantation in the Superficial Femoral Artery on Angioscopy. *Circ J*. 2016;80:2249-51.

11. Ishihara T, **Iida O**, Okamoto S, Fujita M, Masuda M, Nanto K, Shiraki T, Kanda T, Tsujimura T, Okuno S, Yanaka K, Uematsu M. Potential mechanisms of in-stent occlusion in the femoropopliteal artery: an angioscopic assessment. *Cardiovasc Interv Ther*. 2016 Jul 18. [Epub ahead of print]
12. Tomoi Y, Soga Y, Fujihara M, **Iida O**, Shintani Y, Zen K, Ando K. Outcomes of Endovascular Therapy for Upper Extremity Peripheral Artery Disease With Critical Hand Ischemia. Tomoi Y, Soga Y, Fujihara M, Iida O, Shintani Y, Zen K, Ando K. *J Endovasc Ther*. 2016;23:717-22.
13. Utsunomiya M, **Iida O**, Yamauchi Y, Nakano M, Soga Y, Kawasaki D, Takahara M, Nakamura M. Influence of Repeat Intervention on the Risk of Major Amputation After Infrapopliteal Angioplasty for Critical Limb Ischemia. *J Endovasc Ther*. 2016;23:710-6.
14. Okuno S, **Iida O**, Shiraki T, Fujita M, Masuda M, Okamoto S, Ishihara T, Nanto K, Kanda T, Takahara M, Uematsu M. Impact of Calcification on Clinical Outcomes After Endovascular Therapy for Superficial Femoral Artery Disease: Assessment Using the Peripheral Artery Calcification Scoring System. *J Endovasc Ther*. 2016;23:731-7.
15. Shiraki T, **Iida O**, Suemitsu K, Tsuji Y, Uematsu M. Retrograde Approach Using Surgical Cutdown Technique for Limb Salvage in a Case of Critical Limb Ischemia With Severely Calcified Tibial Occlusive Disease. *Vasc Endovascular Surg*. 2016;50:295-8.
16. Masuda M, Fujita M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Kanda T, Shiraki T, Sunaga A, Matsuda Y, Uematsu M. Steerable versus non-steerable sheaths during pulmonary vein isolation: impact of left atrial enlargement on the catheter-tissue contact force. *J Interv Card Electrophysiol*. 2016;47:99-107.
17. Murata N, Takahara M, Soga Y, Nakano M, Yamauchi Y, Zen K, Kawasaki D, Yokoi H, Tosaka A, Tanaka N, **Iida O**. Drug-Eluting Stent vs Percutaneous Transluminal Angioplasty for Treatment of Femoropopliteal In-Stent Restenosis: Results From a Retrospective 1-Year Multicenter Study. *J Endovasc Ther*. 2016;23:642-7.
18. Tomoi Y, Soga Y, **Iida O**, Shiraki T, Kobayashi Y, Hiramori S, Ando K. Impact of Drug-Eluting Stent Implantation for Femoropopliteal In-Stent Occlusion. *J Endovasc Ther*. 2016;23:461-7.
19. Suzuki K, Mizutani Y, Soga Y, **Iida O**, Kawasaki D, Yamauchi Y, Hirano K, Koshida R, Kamoi D, Tazaki J, Higashitani M, Shintani Y, Yamaoka T, Okazaki S, Suematsu N, Tsuchiya T, Miyashita Y, Shinozaki N, Takahashi H, Inoue N. Efficacy and Safety of Endovascular Therapy for Aortoiliac TASC D Lesions. *Angiology*. 2017;68:67-73.
20. Okamoto S, **Iida O**, Takahara M, Yamauchi Y, Hirano K, Soga Y, Suzuki K, Uematsu M. Impact of Perioperative Complications After Endovascular Therapy in Diabetic Patients With Critical Limb Ischemia due to Isolated Infrapopliteal Lesions. *J Endovasc Ther*. 2016;23:371-7.

21. Tsuchiya T, **Iida O**, Shiraki T, Soga Y, Hirano K, Suzuki K, Yamaoka T, Miyashita Y, Kitayama M, Kajinami K. Clinical characteristics of patients with Rutherford category IV, compared with V and VI. *SAGE Open Med.* 2015;3:2050312115597087.
22. Soga Y, Takahara M, **Iida O**, Nakano M, Yamauchi Y, Zen K, Kawasaki D, Ando K. Propensity Score Analysis Comparing Clinical Outcomes of Drug-Eluting vs Bare Nitinol Stents in Femoropopliteal Lesions. *J Endovasc Ther.* 2016;23:33-9.
23. Masuda M, Fujita M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Kanda T, Shiraki T, Sunaga A, Matsuda Y, Uematsu M. Influence of underlying substrate on atrial tachyarrhythmias after pulmonary vein isolation. *Heart Rhythm.* 2016;13:870-8.
24. Hioki H, Miyashita Y, Shiraki T, **Iida O**, Uematsu M, Miura T, Ebisawa S, Ikeda U. Impact of deteriorated calcium-phosphate homeostasis on amputation-free survival after endovascular revascularization in patients with critical limb ischemia on hemodialysis. *Vasc Med.* 2016;21:137-43.
25. Higashimori A, **Iida O**, Yamauchi Y, Kawasaki D, Nakamura M, Soga Y, Zen K, Yokoi Y; OLIVE Investigators. Outcomes of One straight-line flow with and without pedal arch in patients with critical limb ischemia. *Catheter Cardiovasc Interv.* 2016;87:129-33.
26. Soga Y, Takahara M, **Iida O**, Mii S, Okazaki J, Nakano M, Yamauchi Y, Ando K. Bypass Surgery vs. Drug-Eluting Stent for Trans-Atlantic Inter-Society Consensus-II (TASCII) C or D Femoropopliteal Lesions. *Circ J.* 2015;79:2688-95.
27. Soga Y, Takahara M, **Iida O**, Suzuki K, Hirano K, Kawasaki D, Shintani Y, Yamaoka T, Ando K. Relationship Between Primary Patency and Lesion Length Following Bare Nitinol Stent Placement for Femoropopliteal Disease. *J Endovasc Ther.* 2015;22:862-7.
28. Kanda T, Masuda M, Sunaga A, Fujita M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Shiraki T, Sera F, Uematsu M. Fabry cardiomyopathy presenting with a high defibrillation threshold: A short case report. *J Arrhythm.* 2015;31:170-1.
29. Kanda T, Fujita M, **Iida O**, Masuda M, Okamoto S, Ishihara T, Nanto K, Shiraki T, Takahara M, Sakata Y, Uematsu M. Novel Echocardiographic Approach to the Accurate Measurement of Pulmonary Vascular Resistance Based on a Theoretical Formula in Patients With Left Heart Failure - Pilot Study. *Circ J.* 2015;79:2408-13.
30. Shiraki T, **Iida O**, Takahara M, Soga Y, Mii S, Okazaki J, Kuma S, Yamaoka T, Kamoi D, Shintani Y, Ishikawa T, Kitano I, Uematsu M. Predictors of 2-Year Mortality and Risk Stratification After Surgical or Endovascular Revascularization of Infringuinal Artery Disease in Hemodialysis Patients With Critical Limb Ischemia. *J Endovasc Ther.* 2015;22:719-24.
31. Nakano M, Hirano K, Yamauchi Y, **Iida O**, Soga Y, Kawasaki D, Yamaoka T, Suematsu N, Suzuki K. Three-year clinical outcome after infrapopliteal angioplasty for critical limb ischemia in hemodialysis patients with minor or major tissue loss. *Catheter Cardiovasc Interv.* 2015;86:289-98.

32. Shiraki T, **Iida O**, Okamoto S, Ishihara T, Fujita M, Uematsu M. Usefulness of Endovascular Therapy for Penile Gangrene Secondary to Calciphylaxis: A Case Report. *Ann Vasc Surg.* 2015;29:1451.e11-5.
33. Ishihara T, Fujita M, **Iida O**, Tsujimura T, Uematsu M. Rupture of Calcified Nodule 105 Months After Sirolimus-Eluting Stent Implantation Observed on Coronary Angioscopy and Optical Frequency Domain Imaging In Vivo. *Circ J.* 2015;79:2278-9.
34. Nakano M, Hirano K, **Iida O**, Yamauchi Y, Soga Y, Kawasaki D, Tazaki J, Suzuki K, Fujiwara M, Yamaoka T. Clinical efficacy of infrapopliteal endovascular procedures for hemodialysis patients with critical limb ischemia. *Ann Vasc Surg.* 2015;29:1225-34.
35. Sunaga A, Masuda M, Kanda T, Fujita M, **Iida O**, Okamoto S, Ishihara T, Matsuda Y, Watanabe T, Sakata Y, Uematsu M. A low fibrillatory wave amplitude predicts sinus node dysfunction after catheter ablation in patients with persistent atrial fibrillation. *J Interv Card Electrophysiol.* 2015;43:253-61.
36. Kawasaki D, **Iida O**, Fukunaga M, Kato M, Ohkubo N. Wire Passages of 0.035-inch Looped Wire Technique for Femoropopliteal Long Total Occlusions. *J Atheroscler Thromb.* 2015;22:1071-9.
37. Kanda T, Masuda M, Sunaga A, Fujita M, **Iida O**, Okamoto S, Ishihara T, Watanabe T, Takahara M, Sakata Y, Uematsu M. Low left atrial appendage flow velocity predicts recurrence of atrial fibrillation after catheter ablation of persistent atrial fibrillation. *J Cardiol.* 2015;66:377-81.
38. Kobayashi Y, Sakaki M, Yasuoka T, **Iida O**, Dohi T, Uematsu M. Endovascular repair with contralateral external-to-internal iliac artery bypass grafting: a case series. *BMC Res Notes.* 2015;8:183.
39. Soga Y, **Iida O**, Takahaera M, Hirano K, Suzuki K, Kawasaki D. Beta-blocker Treatment Does Not Worsen Critical Limb Ischemia in Patients Receiving Endovascular Therapy. *J Atheroscler Thromb.* 2015;22:481-9.
40. Sato K, **Iida O**, Takahara M, Soga Y, Suzuki K, Tanigawa T, Ito M, Uematsu M. Effect of perioperative complications after endovascular therapy in patients with peripheral artery disease due to femoropopliteal lesions. *J Vasc Surg.* 2015;61:1272-7.
41. Shiraki T, **Iida O**, Takahara M, Soga Y, Yamauchi Y, Hirano K, Kawasaki D, Fujihara M, Utsunomiya M, Tazaki J, Yamaoka T, Shintani Y, Suematsu N, Suzuki K, Miyashita Y, Tsuchiya T, Uematsu M. Predictors of Delayed Wound Healing after Endovascular Therapy of Isolated Infrapopliteal Lesions Underlying Critical Limb Ischemia in Patients with High Prevalence of Diabetes Mellitus and Hemodialysis. *Eur J Vasc Endovasc Surg.* 2015;49:565-73.
42. Suematsu N, **Iida O**, Takahara M, Yamauchi Y, Soga Y, Nakano M, Hirano K, Kawasaki D, Yamaoka T, Suzuki K, Shintani Y, Miyashita Y, Tazaki J, Meno H, Inou T. Prognostic Factors

in Hemodialysis Patients Undergoing Endovascular Treatment for Critical Limb Ischemia due to Isolated Below-the-Knee Disease. *J Atheroscler Thromb.* 2015;22:404-14.

43. Murata N, Soga Y, **Iida O**, Yamauchi Y, Hirano K, Kawasaki D, Fujihara M, Tomoi Y. Complex relationship of body mass index with mortality in patients with critical limb ischemia undergoing endovascular treatment. *Eur J Vasc Endovasc Surg.* 2015;49:297-305.
44. Takahara M, **Iida O**, Soga Y, Kodama A, Azuma N; SPINACH study investigators. Absence of Preceding Intermittent Claudication and its Associated Clinical Features in Patients with Critical Limb Ischemia. *J Atheroscler Thromb.* 2015;22:718-25.
45. Okamoto S, **Iida O**, Nakamura M, Yamauchi Y, Fukunaga M, Yokoi Y, Soga Y, Zen K, Hirano K, Suematsu N, Suzuki K, Shintani Y, Miyashita Y, Urasawa K, Kitano I, Yamaoka T, Ohura N, Hamasaki T, Uematsu M, Nanto S; on behalf of the OLIVE Investigators. Postprocedural Skin Perfusion Pressure Correlates With Clinical Outcomes 1 Year After Endovascular Therapy for Patients With Critical Limb Ischemia. *Angiology.* 2015;66:862-6.
46. Shintani Y, Soga Y, Takahara M, **Iida O**, Kawasaki D, Yamauchi Y, Suzuki K, Hirano K, Kawasaki T. Clinical Outcomes of SMART Versus Luminexx Nitinol Stent Implantation for Aortoiliac Artery Disease: A Propensity Score-Matched Multicenter Study. *Angiology.* 2015;66:875-81.
47. Soga Y, **Iida O**, Takahara M, Hirano K, Suzuki K, Kawasaki D, Miyashita Y, Tsuchiya T. Two-year life expectancy in patients with critical limb ischemia. *JACC Cardiovasc Interv.* 2014;7:1444-9.
48. Nanto K, **Iida O**, Takahara M, Soga Y, Suzuki K, Hirano K, Kawasaki D, Shintani Y, Suematsu N, Yamaoka T, Uematsu M. Effect of Cilostazol Following Endovascular Intervention for Peripheral Artery Disease. *Angiology.* 2015;66:774-8.
49. Ishihara T, **Iida O**, Awata M, Nanto K, Shiraki T, Okamoto S, Iida T, Fujita M, Watanabe T, Nanto S, Uematsu M. Extensive arterial repair one year after paclitaxel-coated nitinol drug-eluting stent vs. bare-metal stent implantation in the superficial femoral artery. *Cardiovasc Interv Ther.* 2015;30:51-6.
50. Miura T, Soga Y, Miyashita Y, **Iida O**, Kawasaki D, Hirano K, Suzuki K, Ikeda U. Five-year prognosis after endovascular therapy in claudicant patients with iliofemoral artery disease. *J Endovasc Ther.* 2014;21:381-8.
51. Soga Y, Mii S, **Iida O**, Okazaki J, Kuma S, Hirano K, Suzuki K, Kawasaki D, Yamaoka T, Kamoi D, Shintani Y. Propensity Score Analysis of Clinical Outcome After Bypass Surgery vs. Endovascular Therapy for Infringuinal Artery Disease in Patients With Critical Limb Ischemia. *J Endovasc Ther.* 2014;21:243-53.
52. Shiraki T, **Iida O**, Takahara M, Okamoto S, Kitano I, Tsuji Y, Terashi H, Uematsu M. Predictive scoring model of mortality after surgical or endovascular revascularization in patients with critical limb ischemia. *J Vasc Surg.* 2014;60:383-9.

53. Soga Y, **Iida O**, Suzuki K, Hirano K, Kawasaki D, Shintani Y, Suematsu N, Yamaoka T, Tsuchiya T, Miyashita Y, Okazaki S, Shinozaki N, Takahashi H. Clinical Impact of Bisoprolol Versus Carvedilol in Patients Undergoing Femoropopliteal Stenting. *J Atheroscler Thromb*. 2014;21:691-702.
54. Aihara H, Soga Y, **Iida O**, Suzuki K, Tazaki J, Shintani Y, Miyashita Y; REAL-AI Registry Investigators. Long-Term Outcomes of Endovascular Therapy for Aortoiliac Bifurcation Lesions in the Real-AI Registry. *J Endovasc Ther*. 2014;21:25-33.
55. Azuma N, **Iida O**, Takahara M, Soga Y, Kodama A. Surgical reconstruction versus peripheral intervention in patients with critical limb ischemia - a prospective multicenter registry in Japan: The SPINACH study design and rationale. *Vascular*. 2014;22:411-20.
56. Iida T, **Iida O**, Okamoto S, Dohi T, Nanto K, Uematsu M. Endovascular therapy with novel high anterior tibial artery puncture for limb salvage in a case of critical lower limb ischemia. *Cardiovasc Interv Ther*. 2014;29:363-7.
57. Dohi T, **Iida O**, Soga Y, Hirano K, Suzuki K, Takahara M, Uematsu M, Nanto S. Incidence, predictors, and prognosis of in-stent occlusion after endovascular treatment with nitinol stents for femoropopliteal lesions. *J Vasc Surg*. 2014;59:1009-1015.e1.
58. Ishihara T, Awata M, Fujita M, Watanabe T, **Iida O**, Ishida Y, Nanto S, Uematsu M. Angioscopic Assessment of Peri-Stent Contrast Staining Following Drug-Eluting Stent Implantation. *Circ J*. 2013;78:122-7.
59. Mizoguchi H, **Iida O**, Dohi T, Tomoda K, Kimura H, Inoue K, Iwata T, Tei K, Miura T. Abdominal Aortic Aneurysmal and Endovascular Device Infection With Iliopsoas Abscess Caused by Mycobacterium Bovis as a Complication of Intravesical Bacillus Calmette-Guérin Therapy. *Ann Vasc Surg*. 2013;27:1186.e1-5.
60. Suzuki K, **Iida O**, Yamauchi Y, Nakano M, Soga Y, Kawasaki D, Tazaki J, Yamaoka T, Suematsu N, Shintani Y, Miyashita Y, Inoue N, Meguro T. Impact of Diabetes Mellitus on Critical Limb Ischemia With Below the Knee Disease: Japan Below-the-Knee Artery Treatment Subanalysis. *Angiology*. 2013 Aug 21. [Epub ahead of print]
61. Soga Y, **Iida O**, Suzuki K, Hirano K, Kawasaki D, Shintani Y, Suematsu N, Yamaoka T. Initial and 3-year results after subintimal versus intraluminal approach for long femoropopliteal occlusion treated with a self-expandable nitinol stent. *J Vasc Surg*. 2013;58:1547-55.
62. Ishihara T, **Iida O**, Awata M, Shiraki T, Nanto S, Uematsu M. Angioscopically apparent large thrombus and uncovered stent struts 6 months after late stent thrombosis of a paclitaxel-coated nitinol drug-eluting stent implanted in the superficial femoral artery. *Cardiovasc Interv Ther*. 2014;29:82-5.
63. Watanabe T, Fujita M, Awata M, **Iida O**, Okamoto S, Ishihara T, Uematsu M. Integrity of stent polymer layer after drug-eluting stent implantation: in vivo comparison of sirolimus-, paclitaxel-, zotarolimus- and everolimus-eluting stents. *Cardiovasc Interv Ther*. 2014;29:4-10.

64. Tomoi Y, Soga Y, **Iida O**, Hirano K, Suzuki K, Kawasaki D, Yamauchi Y, Miyashita Y, Tazaki J, Nobuyoshi M. Efficacy of statin treatment after endovascular therapy for isolated below-the-knee disease in patients with critical limb ischemia. *Cardiovasc Interv Ther.* 2013;28:374-82.
65. Takahara M, **Iida O**, Soga Y, Hirano K, Yamaoka T, Kawasaki D, Suzuki K, Suematsu N, Shintani Y, Miyashita Y, Kaneto H, Shimomura I. Seasonal Variation in Critical Limb Ischemia Requiring Endovascular Therapy: An Analysis of a Multicenter Database of Japanese Patients with Critical Limb Ischemia Undergoing Endovascular Therapy. *J Atheroscler Thromb.* 2013;20:726-32.
66. Ishihara T, Awata M, Fujita M, Watanabe T, **Iida O**, Ishida Y, Nanto S, Uematsu M. Very late stent thrombosis 5 years after implantation of a sirolimus-eluting stent observed by angioscopy and optical coherence tomography. *JACC Cardiovasc Interv.* 2013;6:e28-30.
67. Ishihara T, **Iida O**, Awata M, Nanto K, Nanto S, Uematsu M. Angioscopic Assessment of Early Phase Arterial Repair After Paclitaxel-Coated Nitinol Drug-Eluting Stent Implantation in the Superficial Femoral Artery. *Circ J.* 2013;77:1838-43.
68. Dohi T, **Iida O**, Okamoto S, Nanto K, Nanto S, Uematsu M. Mid-term clinical outcome following endovascular therapy in patients with chronic aortic occlusion. *Cardiovasc Interv Ther.* 2013;28:327-32.
69. Ishihara T, **Iida O**, Awata M, Fujita M, Watanabe T, Ishida Y, Nanto S, Uematsu M. Two-Month Angioscopic Evaluation of Superficial Femoral Artery Treated With Paclitaxel-Coated Nitinol Drug-Eluting Stent. *Circ J.* 2013;77:1880-2.
70. Sakamoto Y, Hirano K, **Iida O**, Soga Y, Suzuki K, Muramatsu T, Tsukahara R. Five-year outcomes of self-expanding nitinol stent implantation for chronic total occlusion of the superficial femoral and proximal popliteal artery. *Catheter Cardiovasc Interv.* 2013;82:E251-6.
71. Takahara M, Kaneto H, Katakami N, **Iida O**, Matsuoka TA, Shimomura I. Effect of sarpogrelate treatment on the prognosis after endovascular therapy for critical limb ischemia. *Heart Vessels.* 2014;29:563-7.
72. Ishihara T, Awata M, Sera F, Fujita M, Watanabe T, **Iida O**, Ishida Y, Nanto S, Uematsu M. Arterial Repair 4 Months After Zotarolimus-Eluting Stent Implantation Observed on Angioscopy. *Circ J.* 2013;77:1186-92.
73. Nakano M, Hirano K, **Iida O**, Soga Y, Kawasaki D, Suzuki K, Miyashita Y. Prognosis of Critical Limb Ischemia in Hemodialysis Patients After Isolated Infrapopliteal Balloon Angioplasty: Results From the Japan Below-the-Knee Artery Treatment (J-BEAT) Registry. *J Endovasc Ther.* 2013;20:113-24.
74. Soga Y, Tomoi Y, Sato K, **Iida O**, Yokoi H. Clinical outcome after endovascular treatment for isolated common femoral and popliteal artery disease. *Cardiovasc Interv Ther.* 2013;28:250-7.

75. Soga Y, **Iida O**, Kawasaki D, Hirano K, Yamaoka T, Suzuki K. Impact of Cilostazol on Angiographic Restenosis after Balloon Angioplasty for Infrapopliteal Artery Disease in Patients with Critical Limb Ischemia. *Eur J Vasc Endovasc Surg.* 2012;44:577-81.
76. Soga Y, **Iida O**, Kawasaki D, Yamauchi Y, Suzuki K, Hirano K, Koshida R, Kamoi D, Tazaki J, Higashitani M, Shintani Y, Yamaoka T, Okazaki S, Suematsu N, Tsuchiya T, Miyashita Y, Shinozaki N, Takahashi H; on behalf of REAL-AI investigators. Contemporary Outcomes After Endovascular Treatment for Aorto-Iliac Artery Disease. *Circ J.* 2012;76:2697-704.
77. Ishihara T, **Iida O**, Tosaka A, Soga Y, Sakamoto Y, Hirano K, Nanto S, Uematsu M. Severity of Coronary Artery Disease Affects Prognosis of Patients With Peripheral Artery Disease. *Angiology.* 2013;64:417-22.
78. Takahara M, Kaneto H, **Iida O**, Katakami N, Matsuoka TA, Ikeda M, Shimomura I. Association of diabetes and hemodialysis with ankle pressure and ankle-brachial index in Japanese patients with critical limb ischemia. *Diabetes Care.* 2012;35:2000-4.
79. Ishihara T, **Iida O**, Okamoto S, Dohi T, Sato K, Nanto K, Fujita M, Watanabe T, Awata M, Sera F, Tanaka N, Ishida Y, Nanto S, Uematsu M. Successful outside-the-stent stenting for in-stent chronic total occlusion in the common iliac artery. *Cardiovasc Interv Ther.* 2012;27:131-6.
80. Tosaka A, Soga Y, **Iida O**, Ishihara T, Hirano K, Suzuki K, Yokoi H, Nanto S, Nobuyoshi M. Classification and clinical impact of restenosis after femoropopliteal stenting. *J Am Coll Cardiol.* 2012;59:16-23.
81. Takahara M, Kaneto H, **Iida O**, Katakami N, Sakamoto F, Matsuoka TA, Ikeda M, Shimomura I. No association of diabetic duration or insulin use with the prognosis of critical limb ischemia after endovascular therapy. *J Atheroscler Thromb.* 2011;18:1102-9.
82. Soga Y, **Iida O**, Hirano K, Suzuki K, Tosaka A, Yokoi H, Nobuyoshi M. Utility of new classification based on clinical and lesional factors after self-expandable nitinol stenting in the superficial femoral artery. *J Vasc Surg.* 2011;54:1058-66.
83. Soga Y, **Iida O**, Hirano K, Suzuki K, Kawasaki D, Miyashita Y, Tsuchiya T, Nobuyoshi M. Impact of cilostazol after endovascular treatment for infrainguinal disease in patients with critical limb ischemia. *J Vasc Surg.* 2011;54:1659-67.
84. Soga Y, **Iida O**, Hirano K, Suzuki K, Yokoi H, Nobuyoshi M. Restenosis after stent implantation for superficial femoral artery disease in patients treated with cilostazol. *Catheter Cardiovasc Interv.* 2012;79:541-8.
85. Tsuji Y, Kitano I, **Iida O**, Kajita S, Sawada K, Nanto S. Popliteal pseudoaneurysm caused by stent fracture. *Ann Vasc Surg.* 2011;25:840.e5-8.
86. Awata M, Uematsu M, Sera F, Ishihara T, Watanabe T, Fujita M, Onishi T, **Iida O**, Ishida Y, Nanto S, Nagata S. Angioscopic assessment of arterial repair following biodegradable polymer-coated biolimus A9-eluting stent implantation. - Comparison with durable polymer-coated sirolimus-eluting stent-. *Circ J.* 2011;75:1113-9.

87. Nanto S, Sakata Y, **Iida O**. Evidence of coronary intervention in Japan. *Nihon Rinsho*. 2011;69:217-23. Review. Japanese.
88. Bito Y, Sakaki M, **Iida O**, Inoue K, Yoshioka Y, Mizoguchi H. Clinical management of lower limb ischemia secondary to a persistent sciatic artery aneurysm: report of a case. *Surg Today*. 2011;41:402-5.
89. Suzuki K, **Iida O**, Soga Y, Hirano K, Inoue N, Uematsu M, Yokoi H, Muramatsu T, Nanto S, Nobuyoshi M, Meguro T. Long-term results of the S.M.A.R.T. Control(TM) stent for superficial femoral artery lesions, J-SMART registry. *Circ J*. 2011;75:939-44.
90. Terashi H, Iwayama T, **Iida O**, Kitano I, Tsuji Y. Dynamic skin perfusion pressure: a new measure of assessment for wound healing capacity and alternative angiosome in critical limb ischemia. *Plast Reconstr Surg*. 2010;126:215e-218e.
91. Takahara M, Kaneto H, **Iida O**, Gorogawa S, Katakami N, Matsuoka TA, Ikeda M, Shimomura I. The influence of glycemic control on the prognosis of Japanese patients undergoing percutaneous transluminal angioplasty for critical limb ischemia. *Diabetes Care*. 2010;33:2538-42.
92. Takahara M, Kaneto H, **Iida O**, Gorogawa S, Ikeda M. High prevalence of glucose intolerance in Japanese patients with peripheral arterial disease. *Diabetes Res Clin Pract*. 2011;91:e24-5.
93. Onishi T, Uematsu M, Watanabe T, Fujita M, Awata M, **Iida O**, Sera F, Hirano Y, Nanto S, Nagata S. Objective interpretation of dobutamine stress echocardiography by diastolic dyssynchrony imaging: a practical approach. *J Am Soc Echocardiogr*. 2010 ;23:1103-8.
94. Soga Y, **Iida O**, Hirano K, Yokoi H, Nanto S, Nobuyoshi M. Mid-term clinical outcome and predictors of vessel patency after femoropopliteal stenting with self-expandable nitinol stent. *J Vasc Surg*. 2010;52:608-15.
95. Miyashita Y, Saito S, Miyamoto A, **Iida O**, Nanto S. Cilostazol increases skin perfusion pressure in severely ischemic limbs. *Angiology*. 2011;62:15-7.
96. Awata M, Nanto S, Uematsu M, Morozumi T, Watanabe T, Onishi T, **Iida O**, Sera F, Minamiguchi H, Kotani J, Nagata S. Heterogeneous arterial healing in patients following paclitaxel-eluting stent implantation: comparison with sirolimus-eluting stents. *JACC Cardiovasc Interv*. 2009;2:453-8.
97. Onishi T, Uematsu M, Nanto S, Morozumi T, Watanabe T, Awata M, **Iida O**, Sera F, Nagata S. Detection of diastolic abnormality by dyssynchrony imaging: correlation with coronary artery disease in patients presenting with visibly normal wall motion. *Circ J*. 2009;73:125-31.
98. Awata M, Nanto S, Uematsu M, Morozumi T, Watanabe T, Onishi T, **Iida O**, Sera F, Kotani J, Hori M, Nagata S. Angioscopic comparison of neointimal coverage between zotarolimus- and sirolimus-eluting stents. *J Am Coll Cardiol*. 2008;52:789-90.
99. Awata M, Kotani J, Uematsu M, Morozumi T, Watanabe T, Onishi T, **Iida O**, Sera F, Nanto S, Hori M, Nagata S. Serial angioscopic evidence of incomplete neointimal coverage after

sirolimus-eluting stent implantation: comparison with bare-metal stents. *Circulation*. 2007;116:910-6.

100. Goto Y, Awata M, Uematsu M, Yano M, Morozumi T, Kotani J, Watanabe T, Onishi T, **Iida O**, Sera F, Nanto S, Nagata S. Shoshin beriberi complicating severe pulmonary hypertension: a case report. *J Cardiol*. 2007;49:361-5. Japanese.
101. Onishi T, Uematsu M, Nanto S, **Iida O**, Morozumi T, Kotani J, Awata M, Nagata S. Positive isovolumic relaxation velocity detected by a spectral tissue Doppler mapping technique as an indicator of coronary artery disease: a prospective study. *J Am Soc Echocardiogr*. 2007;20:158-64.

Abstract

1. Long-term Clinical Efficacy of Endovascular Therapy for Patients with Critical Limb Ischemia due to Pure Isolated Infrapopliteal Lesions. *Circulation* 2012; (Abstract 13627)
2. Angiographic Restenosis and Its Clinical Impact after Below-the-knee Angioplasty. *J Am Coll Cardiol* 2012; 61th ACC (Abstract E2087)
3. Long-term Clinical Outcomes after Endovascular Therapy for Patients with Critical Limb Ischemia due to Isolated Below the Knee Lesions. *Circulation* 2011; (Abstract 11141)
4. Impact of Angiosome in Endovascular Therapy (EVT) on the Limb Salvage for the Patients with Critical Limb Ischemia (CLI) Presenting with Isolated Infrapopliteal Lesions. *J Am Coll Cardiol* 2011; 60th ACC (Abstract E1667)
5. Time Course of the Restenosis Following Nitinol Stenting in the Superficial Femoral Artery and the Factors Associated with Early and Late Restenoses. *Circulation* 2010 (Abstract 12185)
6. Stent Fracture, TASC II CD lesion as Restenosis Factors, and Cilostazol as a Negative-restenosis Factor Within a Year Following Nitinol Stent Implantation in the Superficial Femoral Artery. *J Am Coll Cardiol* 2010; 59th ACC (Abstract .E1688)
7. Acquiring Direct Flow Based on the Angiosome Concept in Endovascular Therapy Saves More Limbs than Increasing the Number of Patent Run-off Vessels in Patients with Critical Limb Ischemia. *Circulation* 2009 (Abstract 4427)
8. Factors Differ Between Early and Late Restenosis Following Nitinol Stent Implantation in the Superficial Femoral Artery. *Circulation* 2009 (Abstract 4424)
9. Stent Fracture does not Affect Long Term Patency in the Femoro-popliteal Artery: Four Years Experience. *Circulation* 2008 (Abstract 6189)
10. Low Ejection Fraction and Poor Performance Status Negate the Beneficial Effects of Complete Revascularization in Patients with Critical Limb Ischemia. *J Am Coll Cardiol* 2008; 57th (Abstract)
11. Complete Revascularization to the Pedal Arch may Improve the Survival of Patient with Critical Limb Ischemia. *Circulation* 2007 (Abstract 3228)
12. Cilostazol Attenuates Angiographic Restenosis in Patients with Femoro-popliteal Lesions *Circulation* 2007 (Abstract 3224)
13. Morphology of the Stent Fracture Influences the Outcome Following Superficial Femoral Artery Stenting. *J Am Coll Cardiol* 2007; 56th (Abstract)
14. Successful Revascularization of at Least a Branch of the Tibial Artery to the Pedal Arch Contributes Limb Salvage in Critical Limb Ischemia. *J Am Coll Cardiol* 2007; 56th (Abstract)

Curriculum Vitae - Osamu Iida

15. Cilostazol Reduces Target Lesion Revascularization in the Femoro-Popliteal Lesions in Patients Complaining of Claudication. *Circulation* 2005 (Abstract 2242)
16. Exercise Adversely Affects Stent Fracture in the Superficial Femoral Artery. *Circulation* 2005 (Abstract 2241)
17. Stenosis in the Coronary Artery of Normally Contracting Heart: Detection by a Simplified tissue Doppler Mapping Technique. *J Am Coll Cardiol* 2004; 53th: 371A (Abstract)