

Publications - First Author

---

1. **Iida O**, Ohki T, Soga Y, Suematsu N, Nakama T, Yamaoka T, Tobita K, Ichihashi S. Twelve-Month Outcomes From the Japanese Post-Market Surveillance Study of the Viabahn Endoprosthesis as Treatment for Symptomatic Peripheral Arterial Disease in the Superficial Femoral Arteries. *J Endovasc Ther.* 2021 Dec 15:15266028211067739. doi: 10.1177/15266028211067739. Online ahead of print.
2. **Iida O**, Takahara M, Soga Y, Fujihara M, Kawasaki D, Hirano K, Choi D, Mano T. Vessel Diameter Evaluated by Intravascular Ultrasound Versus Angiography. *J Endovasc Ther.* 2021 Sep 27:15266028211047946. doi: 10.1177/15266028211047946. Online ahead of print.
3. **Iida O**, Urasawa K, Shibata Y, Yamamoto Y, Ando H, Fujihara M, Nakama T, Miyashita Y, Mori S, Diaz-Cartelle J, Soga Y. Clinical Safety and Efficacy of Rotational Atherectomy in Japanese Patients with Peripheral Arterial Disease Presenting Femoropopliteal Lesions: The J-SUPREME and J-SUPREME II Trials. *J Endovasc Ther.* 2021 Sep 13:15266028211045700. doi: 10.1177/15266028211045700. Online ahead of print.
4. **Iida O**, Fujihara M, Kawasaki D, Mori S, Yokoi H, Miyamoto A, Kichikawa K, Nakamura M, Ohki T, Diaz-Cartelle J, Müller-Hülsbeck S, Gray WA, Soga Y. 24-Month Efficacy and Safety Results from Japanese Patients in the IMPERIAL Randomized Study of the Eluvia Drug-Eluting Stent and the Zilver PTX Drug-Coated Stent. *Cardiovasc Intervent Radiol.* 2021;44:1367-1374.
5. **Iida O**, Takahara M, Mano T. Evidence-Experience Gap and Future Perspective on the Treatment of Peripheral Artery Disease. *J Atheroscler Thromb.* 2021;28:1251-1259.
6. **Iida O**, Takahara M, Soga Y, Yamaoka T, Nanto S, Kuratani T, Sakata Y, Mano T. One-Year Outcomes of Heparin-Bonded Stent-Graft Therapy for Real-World Femoropopliteal Lesions and the Association of Patency With the Prothrombotic State Based on the Prospective, Observational, Multicenter Viabahn Stent-Graft Placement for Femoropopliteal Diseases Requiring Endovascular Therapy (VANQUISH) Study. *J Endovasc Ther.* 2021;28:123-131.
7. **Iida O**, Takahara M, Soga Y, Fujihara M, Kawasaki D, Hirano K, Choi D, Mano T. A Novel Angiographic Risk Score for Femoropopliteal Interventions. *J Endovasc Ther.* 2020;27:967-973.
8. **Iida O**, Takahara M, Kohsaka S, Soga Y, Fujihara M, Mano T, Shinke T, Amano T, Ikari Y. Impact of Institutional Volume on Critical In-Hospital Complications Adjusted for Patient- and Limb-Related Characteristics: An Analysis of a Nationwide Japanese Registry of Endovascular Interventions for PAD. *J Endovasc Ther.* 2020;27:739-748.
9. **Iida O**, Takahara M, Soga Y, Kodama A, Terashi H, Azuma N; SPINACH investigators. The Association of Preoperative Characteristics with Reintervention Risk in Patients Undergoing Revascularization for Chronic Limb-Threatening Ischemia. *J Atheroscler Thromb.* 2021;28:52-65.
10. **Iida O**, Takahara M, Yamauchi Y, Shintani Y, Sugano T, Yamamoto Y, Kawasaki D, Yokoi H, Miyamoto A, Mano T, Nakamura M; OMOTENASHI investigators. Impact of Hospital

Volume on Clinical Outcomes after Aortoiliac Stenting in Patients with Peripheral Artery Disease. *J Atheroscler Thromb.* 2020;27:516-523.

11. **Iida O**, Takahara M, Mano T. Commentary: Is Calcification Still an Unsolved Issue in the Drug-Solution Era? *J Endovasc Ther.* 2019;26:621-622.
12. **Iida O**, Mano T. Role of IVUS in the Endovascular Treatment of Calcified Femoropopliteal Lesions. *J Endovasc Ther.* 2019;26:331-332.
13. **Iida O**, Urasawa K, Komura Y, Soga Y, Inoue N, Hara H, Yajima J, Nakamura S, Ohki T, Ando H, Hirano K, Horita Y, Kichikawa K, Yokoi Y, Miyamoto A, Nakamura M, Takahara M, Mano T, Nanto S. Self-Expanding Nitinol Stent vs Percutaneous Transluminal Angioplasty in the Treatment of Femoropopliteal Lesions: 3-Year Data From the SM-01 Trial. *J Endovasc Ther.* 2019;26:158-167.
14. **Iida O**, Soga Y, Urasawa K, Saito S, Jaff MR, Wang H, Ookubo H, Yokoi H; MDT-2113 SFA Japan Investigators. Drug-coated balloon versus uncoated percutaneous transluminal angioplasty for the treatment of atherosclerotic lesions in the superficial femoral and proximal popliteal artery: 2-year results of the MDT-2113 SFA Japan randomized trial. *Catheter Cardiovasc Interv.* 2019;93:664-672.
15. **Iida O**, Soga Y, Urasawa K, Saito S, Jaff MR, Wang H, Ookubo H, Yokoi H; MDT-2113 SFA Japan Investigators. Drug-Coated Balloon vs Standard Percutaneous Transluminal Angioplasty for the Treatment of Atherosclerotic Lesions in the Superficial Femoral and Proximal Popliteal Arteries: One-Year Results of the MDT-2113 SFA Japan Randomized Trial. *J Endovasc Ther.* 2018;25:109-117.
16. **Iida O**, Takahara M, Soga Y, Kodama A, Terashi H, Azuma N, ; SPINACH Investigators. Three-Year Outcomes of Surgical Versus Endovascular Revascularization for Critical Limb Ischemia: The SPINACH Study (Surgical Reconstruction Versus Peripheral Intervention in Patients With Critical Limb Ischemia). *Circ Cardiovasc Interv.* 2017;10. pii: e005531. doi: 10.1161/CIRCINTERVENTIONS.117.005531.
17. **Iida O**, Takahara M, Soga Y, Azuma N, Nanto S, Uematsu M; PRIORITY Investigators. Prognostic Impact of Revascularization in Poor-Risk Patients With Critical Limb Ischemia: The PRIORITY Registry (Poor-Risk Patients With and Without Revascularization Therapy for Critical Limb Ischemia). *JACC Cardiovasc Interv.* 2017;10:1147-1157.
18. **Iida O**, Takahara M. What does Intravascular Ultrasound Illustrate? *J Atheroscler Thromb.* 2017;24:473-474.
19. **Iida O**, Takahara M, Mano T. What Does the IN.PACT SFA-Long Study Tell Us? *JACC Cardiovasc Interv.* 2017;10:735-737.

20. **Iida O**, Takahara M, Soga Y, Hirano K, Yamauchi Y, Zen K, Yokoi H, Uematsu M; ZEPHYR investigators. Incidence and its characteristics of repetition of reintervention after drug-eluting stent implantation for femoropopliteal lesion. *J Vasc Surg*. 2016;64:1691-1695.e1.
21. **Iida O**, Takahara M, Soga Y, Hirano K, Yamauchi Y, Zen K, Kawasaki D, Nanto S, Yokoi H, Uematsu M; ZEPHYR Investigators. The Characteristics of In-Stent Restenosis After Drug-Eluting Stent Implantation in Femoropopliteal Lesions and 1-Year Prognosis After Repeat Endovascular Therapy for These Lesions. *JACC Cardiovasc Interv*. 2016;9:828-34.
22. **Iida O**, Nakamura M. Reply: 3-Year Outcomes of the OLIVE Registry, a Prospective Multicenter Study of Patients with Critical Limb Ischemia. *JACC Cardiovasc Interv*. 2016;9:202-3.
23. **Iida O**, Nakamura M, Yamauchi Y, Fukunaga M, Yokoi Y, Yokoi H, Soga Y, Zen K, Suematsu N, Inoue N, Suzuki K, Hirano K, Shintani Y, Miyashita Y, Urasawa K, Kitano I, Tsuchiya T, Kawamoto K, Yamaoka T, Uesugi M, Shinke T, Oba Y, Ohura N, Uematsu M, Takahara M, Hamasaki T, Nanto S; OLIVE Investigators. 3-Year Outcomes of the OLIVE Registry, a Prospective Multicenter Study of Patients With Critical Limb Ischemia: A Prospective, Multi-Center, Three-Year Follow-Up Study on Endovascular Treatment for Infra-Inguinal Vessel in Patients With Critical Limb Ischemia. *JACC Cardiovasc Interv*. 2015;8:1493-502.
24. **Iida O**, Takahara M, Soga Y, Nakano M, Yamauchi Y, Zen K, Kawasaki D, Nanto S, Yokoi H, Uematsu M; ZEPHYR Investigators. 1-Year Results of the ZEPHYR (Zilver PTX for the Femoral Artery and Proximal Popliteal Artery) Registry: Predictors of Restenosis. *JACC Cardiovasc Interv*. 2015;8:1105-1112.
25. **Iida O**, Takahara M, Soga Y, Yamauchi Y, Hirano K, Tazaki J, Yamaoka T, Suematsu N, Suzuki K, Shintani Y, Miyashita Y, Uematsu M. Impact of angiosome-oriented revascularization on clinical outcomes in critical limb ischemia patients without concurrent wound infection and diabetes. *J Endovasc Ther*. 2014;21:607-15.
26. **Iida O**, Takahara M, Soga Y, Suzuki K, Hirano K, Kawasaki D, Shintani Y, Suematsu N, Yamaoka T, Nanto S, Uematsu M. Efficacy of Intravascular Ultrasound in Femoropopliteal Stenting for Peripheral Artery Disease With TASC II Class A to C Lesions. *J Endovasc Ther*. 2014;21:485-492.
27. **Iida O**, Takahara M, Soga Y, Suzuki K, Hirano K, Kawasaki D, Shintani Y, Suematsu N, Yamaoka T, Nanto S, Uematsu M. Shared and Differential Factors Influencing Restenosis Following Endovascular Therapy Between TASC (Trans-Atlantic Inter-Society Consensus) II Class A to C and D Lesions in the Femoropopliteal Artery. *JACC Cardiovasc Interv*. 2014;7:792-8.
28. **Iida O**, Soga Y, Takahara M, Kawasaki D, Yamauchi Y, Suzuki K, Hirano K, Hirano K, Kamoi D, Tazaki J, Higashitani M, Shintani Y, Yamaoka T, Okazaki S, Suematsu N, Tsuchiya T,

Miyashita Y, Shinozaki N, Takahashi H, Uematsu M. Perioperative Complications After Aortoiliac Stenting: Associated Factors and Impact on Follow-up Cardiovascular Prognosis. *Eur J Vasc Endovasc Surg.* 2014;47:131-138.

29. **Iida O**, Takahara M, Soga Y, Yamauchi Y, Hirano K, Tazaki J, Yamaoka T, Suematsu N, Suzuki K, Shintani Y, Miyashita Y, Uematsu M. Worse limb prognosis for indirect vs. direct endovascular revascularization only in patients with critical limb ischemia complicated with wound infection and diabetes mellitus. *Eur J Vasc Endovasc Surg.* 2013;46:575-82.
30. **Iida O**, Soga Y, Takahara M, Kawasaki D, Yamauchi Y, Suzuki K, Hirano K, Ryoji K, Kamoi D, Tazaki J, Higashitani M, Shintani Y, Yamaoka T, Okazaki S, Suematsu N, Tsuchiya T, Miyashita Y, Shinozaki N, Takahashi H, Uematsu M. Efficacy of the S.M.A.R.T. Control vs. other stents for aortoiliac occlusive disease in contemporary clinical practice. *J Endovasc Ther.* 2013;20:431-9.
31. **Iida O**, Yokoi H, Soga Y, Inoue N, Suzuki K, Yokoi Y, Kawasaki D, Zen K, Urasawa K, Shintani Y, Miyamoto A, Hirano K, Miyashita Y, Tsuchiya T, Shinozaki N, Nakamura M, Isshiki T, Hamasaki T, Nanto S; on behalf of the STOP-IC investigators. Cilostazol Reduces Angiographic Restenosis After Endovascular Therapy for Femoropopliteal Lesions in the Sufficient Treatment of Peripheral Intervention by Cilostazol Study. *Circulation.* 2013;127:2307-15.
32. **Iida O**, Soga Y, Yamauchi Y, Hirano K, Kawasaki D, Yamaoka T, Takahara M, Uematsu M. Clinical efficacy of endovascular therapy for patients with critical limb ischemia attributable to pure isolated infrapopliteal lesions. *J Vasc Surg.* 2013;57:974-981.
33. **Iida O**, Nakamura M, Miyamoto A, Kawasaki D, Yokoi Y, Yokoi H, Zen K, Hirano K, Suematsu N, Inoue N, Shintani Y, Miyashita Y, Urasawa K, Kitano I, Yamaoka T, Murakami T, Uesugi M, Tsuchiya T, Shinke T, Oba Y, Ohura N, Hamasaki T, Nanto S on behalf of the OLIVE investigators. Endovascular Treatment for Infringuinal Vessels in Patients with Critical Limb Ischemia: OLIVE Registry, a Prospective, Multicenter Study in Japan with 12-month Follow-up. *Circ Cardiovasc Interv.* 2013;6:68-76.
34. **Iida O**, Soga Y, Kawasaki D, Hirano K, Yamaoka T, Suzuki K, Miyashita Y, Yokoi H, Takahara M, Uematsu M. Angiographic restenosis and its clinical impact after infrapopliteal angioplasty. *Eur J Vasc Endovasc Surg.* 2012;44:425-31.
35. **Iida O**, Soga Y, Hirano K, Kawasaki D, Suzuki K, Miyashita Y, Nanto S, Uematsu M. Anatomical Predictors of Major Adverse Limb Events after Infrapopliteal Angioplasty for Patients with Critical Limb Ischaemia due to Pure Isolated Infrapopliteal Lesions. *Eur J Vasc Endovasc Surg.* 2012;44:318-24.
36. **Iida O**, Soga Y, Hirano K, Kawasaki D, Suzuki K, Miyashita Y, Nanto S, Uematsu M. Midterm Outcomes and Risk Stratification after Endovascular Therapy for Patients with Critical Limb Ischaemia due to Isolated Below-the-knee Lesions. *Eur J Vasc Endovasc Surg.* 2012;43:313-21.

37. **Iida O**, Soga Y, Hirano K, Kawasaki D, Suzuki K, Miyashita Y, Terashi H, Uematsu M. Long-term results of direct and indirect endovascular revascularization based on the angiosome concept in patients with critical limb ischemia presenting with isolated below-the-knee lesions. *J Vasc Surg.* 2012;55:363-370.
38. **Iida O**, Soga Y, Hirano K, Suzuki K, Yokoi H, Nobuyoshi M, Muramatsu T, Inoue N, Nanto S, Uematsu M. Long-term outcomes and risk stratification of patency following nitinol stenting in the femoropopliteal segment: retrospective multicenter analysis. *J Endovasc Ther.* 2011;18:753–761.
39. **Iida O**, Uematsu M, Soga Y, Hirano K, Suzuki K, Yokoi H, Muramatsu T, Inoue N, Nanto S, Nagata S. Timing of the restenosis following nitinol stenting in the superficial femoral artery and the factors associated with early and late restenoses. *Catheter Cardiovasc Interv.* 2011;78:611-7.
40. **Iida O**, Soga Y, Hirano K, Okamoto S, Dohi T, Uematsu M, Yokoi H, Nobuyoshi M, Muramatsu T, Fujita M, Nanto S, Nagata S. Retrospective multicentre analysis of S.M.A.R.T. vs. Luminexx nitinol stent implantation for superficial femoral artery lesions (REAL SL) Registry. 5 years' experience. *Circ J.* 2011;75:421-7.
41. **Iida O**, Nanto S, Uematsu M, Ikeoka K, Okamoto S, Dohi T, Fujita M, Terashi H, Nagata S. Importance of the angiosome concept for endovascular therapy in patients with critical limb ischemia. *Catheter Cardiovasc Interv.* 2010;75:830-6.
42. **Iida O**, Nanto S, Uematsu M, Ikeoka K, Okamoto S, Dohi T, Fujita M, Nagata S. Long-term results of endovascular therapy with nitinol stent implantation for TASC II A/B femoro-popliteal artery lesions: 4 years' experience. *Circ J.* 2009;73:2143-7.
43. **Iida O**, Nanto S, Uematsu M, Ikeoka K, Okamoto S, Nagata S. Influence of stent fracture on the long-term patency in the femoro-popliteal artery: experience of 4 years. *JACC Cardiovasc Interv.* 2009;2:665-71.
44. **Iida O**, Nanto S, Uematsu M, Morozumi T, Ikeoka K, Terashi H, Nagata S. Novel Infra-popliteal Bi-directional Angioplasty in a Patient with Chronic Total Occlusion of the Tibial Artery Complicating Critical Lower Limb Ischemia. *Jpn J Interv Cardiol.* 2008;23:365-9.
45. **Iida O**, Nanto S, Uematsu M, Morozumi T, Kitakaze M, Nagata S. Cilostazol reduces restenosis after endovascular therapy in patients with femoropopliteal lesions. *J Vasc Surg.* 2008;48:144-9.
46. **Iida O**, Nanto S, Uematsu M, Morozumi T, Akahori H, Nagata S. Endovascular therapy for limb salvage in a case of critical lower limb ischemia resulting from fibromuscular dysplasia. *J Vasc Surg.* 2007;46:803-7.

47. **Iida O**, Nanto S, Uematsu M, Morozumi T, Kotani J, Awata M, Onishi T, Ito N, Sera F, Minamiguchi H, Akahori H, Nagata S. Effect of exercise on frequency of stent fracture in the superficial femoral artery. *Am J Cardiol.* 2006;98:272-4.
48. **Iida O**, Nanto S, Uematsu M, Morozumi T, Kotani J, Awata M, Onishi T, Ito N, Oshima F, Minamiguchi H, Kitakaze M, Nagata S. Cilostazol reduces target lesion revascularization after percutaneous transluminal angioplasty in the femoropopliteal artery. *Circ J.* 2005;69:1256-9.



Publications - Co-Author

---

1. Nakao S, **Iida O**, Hata Y, Okamoto S, Ishihara T, Nanto K, Tsujimura T, Takahara M, Mano T. Impact of Cardiac Function on Loss of Patency in Patients With Peripheral Artery Disease Presenting the Femoropopliteal Lesions Endovascularly Treated With New-Generation Devices. *J Endovasc Ther.* 2022 Jan 20;15266028211070963. doi: 10.1177/15266028211070963. Online ahead of print.
2. Kanda T, Masuda M, Asai M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Tsujimura T, Matsuda Y, Hata Y, Uematsu H, Mano T. Extensive Left Atrial Low-Voltage Area During Initial Ablation is Associated with A Poor Clinical Outcome Even Following Multiple Procedures. *J Atr Fibrillation.* 2021;14:20200491. doi: 10.4022/jafib.20200491. eCollection 2021 Aug.
3. Soga Y, Takahara M, **Iida O**, Suzuki K, Mori S, Kawasaki D, Haraguchi K, Yamaoka T, Ando K. Ten-Year Clinical Follow-Up Following Bare-Nitinol Stent Implantation for Femoropopliteal Artery Disease. *J Atheroscler Thromb.* 2021. doi: 10.5551/jat.63225. Online ahead of print.
4. Nakao S, **Iida O**, Ishihara T, Mano T. Five-Year Intra-Arterial Evaluation After Fluoropolymer-Based Drug-Eluting Stent Implantation for a Superficial Femoral Artery. *Circ Rep.* 2021;3:686-687. doi: 10.1253/circrep.CR-21-0089. eCollection 2021 Nov 10.
5. Higashino N, Ishihara T, **Iida O**, Tsujimura T, Hata Y, Toyoshima T, Kurata N, Asai M, Masuda M, Okamoto S, Nanto K, Kanda T, Matsuda Y, Mano T. Early- and Middle-Phase Angioscopic Assessment of Arterial Healing Following Current Drug-Eluting Stent Implantation in Patients With Acute Coronary Syndrome. *Circ Rep.* 2021;3:666-673. doi: 10.1253/circrep.CR-21-0113. eCollection 2021 Nov 10.
6. Kurata N, Masuda M, Kanda T, Asai M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Tsujimura T, Matsuda Y, Hata Y, Uematsu H, Mano T. Left Atrial Localized Low-Voltage Areas Indicate Whole Left Atrial Electrophysiological Degeneration in Atrial Fibrillation Patients. *Circ J.* 2022;86:192-199.
7. Nakao S, Ishihara T, Tsujimura T, **Iida O**, Hata Y, Toyoshima T, Higashino N, Mano T. Effectiveness of hospital lipid-lowering protocol of intensive lipid-lowering therapy for patients with acute coronary syndrome. *J Cardiol.* 2021 Oct 24;S0914-5087(21)00276-8. doi: 10.1016/j.jjcc.2021.10.005. Online ahead of print.
8. Soga Y, **Iida O**, Fujihara M, Kawasaki D, Saito S, Urasawa K, Yokoi H, Fernandez EJ, Guo J, Nakamura M; IN.PACT Japan Post-Market Surveillance Study. Real-World Clinical Outcomes of IN.PACT Admiral Drug-Coated Balloon for Femoropopliteal Artery Disease - 12-Month Results From Japan Post-Market Surveillance Study. *Circ J.* 2021;85:2149-2156.
9. Tomoi Y, Takahara M, Kuramitsu S, Soga Y, **Iida O**, Fujihara M, Kawasaki D, Ando K; IVORY Study Investigators. Subintimal Versus Intraluminal Approach for Femoropopliteal

Chronic Total Occlusions Treated With Intravascular Ultrasound Guidance. *J Am Heart Assoc.* 2021;10:e021903.

10. Soga Y, Fujihara M, Yamamoto Y, Nakamura S, **Iida O**, Kawasaki D, Urasawa K, Ando H, Mori S, Suzuki K, Horie K, Diaz-Cartelle J, Kozuki A. One-year results for Japanese patients in RANGER II SFA. *Heart Vessels.* 2021 Sep 23. doi: 10.1007/s00380-021-01947-3. Online ahead of print.
11. Kodama A, Takahara M, **Iida O**, Soga Y, Mii S, Kitano I, Deguchi J, Fukui D, Komori K, Azuma N; on behalf of the SPINACH investigators. Health Related Quality of Life Over Time After Revascularisation in Patients With Chronic Limb Threatening Ischaemia. *Eur J Vasc Endovasc Surg.* 2021;62:777-785.
12. Takahara M, **Iida O**, Soga Y, Kodama A, Terashi H, Azuma N. Features of asymptomatic contralateral limb in patients with chronic limb-threatening ischemia. *J Cardiol.* 2022;79:153-160.
13. Fujihara M, Takahara M, Yamaoka T, **Iida O**, Kojima T, Tobita K, Nakama T, Kyuragi R, Ichihashi S, Soga Y. Clinical outcomes of endovascular procedure using VIABAHN® VBX covered stent in complex aortoiliac artery disease: Result from AVOCADO study. *Catheter Cardiovasc Interv.* 2021;98:928-937.
14. Matsuda Y, Masuda M, Sakio T, Asai M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Kanda T, Tsujimura T, Hata Y, Uematsu H, Mano T. Safety, efficacy, and cost-performance of a simplified cryoballoon ablation procedure for paroxysmal atrial fibrillation. *J Interv Card Electrophysiol.* 2021 Aug 12. doi: 10.1007/s10840-021-01043-6. Online ahead of print.
15. Masuda M, Asai M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Kanda T, Tsujimura T, Matsuda Y, Hata Y, Uematsu H, Mano T. Low-Voltage-Area Ablation in Paroxysmal Atrial Fibrillation - Extended Follow-up Results of the VOLCANO Trial. *Circ J.* 2022;86:245-252.
16. Ishihara T, Dohi T, Nakamura D, Kikuchi A, Okamoto N, Mori N, **Iida O**, Tsujimura T, Mizote I, Higuchi Y, Yamada T, Nishino M, Mano T, Sakata Y; QUEST Investigators. Impact of in-stent tissue characteristics on excimer laser coronary angioplasty prior to drug-coated balloon treatment. *Int J Cardiol.* 2021;339:28-32.
17. Tsujimura T, **Iida O**, Asai M, Masuda M, Okamoto S, Ishihara T, Nanto K, Kanda T, Matsuda Y, Hata Y, Uematsu H, Toyoshima T, Higashino N, Mano T. Aneurysmal degeneration of fluoropolymer-coated paclitaxel-eluting stent in the superficial femoral artery: a rising concern. *CVIR Endovasc.* 2021;4:56.
18. Ohki T, Kichikawa K, Yokoi H, **Iida O**, Yamaoka T, Maeda K, Kanaoka Y. Long-term results of the Japanese multicenter Viabahn trial of heparin bonded endovascular stent grafts for long and complex lesions in the superficial femoral artery. *J Vasc Surg.* 2021;74:1958-1967.e2.



19. Hata Y, **Iida O**, Ito N, Soga Y, Fukunaga M, Kawasaki D, Fujihara M, Kozuki A, Takahara M, Mano T. Roles of Angioplasty With Drug-Coated Balloon for Chronic Ischemia in Wound Healing. *J Endovasc Ther.* 2021;28:778-787.
20. Kobayashi N, Takahara M, **Iida O**, Soga Y, Kodama A, Hirano K, Nakano M, Yamauchi Y, Komai H, Azuma N. Impact of Dialysis Vintage and Renal Biomarkers on Mortality in Dialysis-Dependent Patients With Critical Limb Ischemia Undergoing Revascularization. *J Endovasc Ther.* 2021;28:716-725.
21. Tsujimura T, Ishihara T, **Iida O**, Hata Y, Toyoshima T, Higashino N, Kurata N, Asai M, Masuda M, Okamoto S, Nanto K, Kanda T, Matsuda Y, Mano T. Arterial Healing 10 Months After Implantation of an Ultrathin-Strut, Biodegradable-Polymer, Sirolimus-Eluting Stent - An Angioscopic Study. *Circ Rep.* 2021;3:316-323.
22. Takahara M, **Iida O**, Soga Y, Kodama A, Terashi H, Azuma N. Impact of homocysteine levels on mortality risk in patients with chronic limb-threatening ischemia undergoing revascularization. *Heart Vessels.* 2021;36:1825-1829.
23. Matsuoka Y, **Iida O**, Suemitsu K, Oka K, Ota N, Izumi M. Use of a fluoropolymer-based paclitaxel-eluting stent for arteriovenous graft outflow vein stenosis in hemodialysis patients. *J Vasc Surg Cases Innov Tech.* 2021;7:326-331. eCollection 2021 Jun.
24. Kodama A, Takahara M, **Iida O**, Soga Y, Terashi H, Kawasaki D, Izumi Y, Mii S, Komori K, Azuma N; SPINACH investigators. Ambulatory Status Over Time after Revascularization in Patients with Chronic Limb-Threatening Ischemia. *J Atheroscler Thromb.* 2021. doi: 10.5551/jat.62892. Online ahead of print.
25. Hata Y, **Iida O**, Takahara M, Asai M, Masuda M, Okamoto S, Ishihara T, Nanto K, Kanda T, Tsujimura T, Matsuda Y, Yanaka K, Mano T. Saphenous vein size as a surrogate marker for mortality of patients with chronic limb-threatening ischemia undergoing endovascular therapy. *J Cardiol.* 2021;78:341-346.
26. Matsuda Y, Masuda M, Asai M, **Iida O**, Kanda T, Mano T. Central retinal artery occlusion after catheter ablation of atrial fibrillation. *Clin Case Rep.* 2021;9:e04255. eCollection 2021 May.
27. Katsuki T, Yamaji K, Soga Y, **Iida O**, Fujihara M, Kawasaki D, Ando K. New lesion after endovascular therapy of femoropopliteal lesions for intermittent claudication. *Catheter Cardiovasc Interv.* 2021;98:E395-E402.
28. Ichihashi S, Takahara M, **Iida O**, Suzuki K, Yamaoka T, Maeda K, Tobita K, Kobayashi T, Nakama T, Shingaki M, Ozaki S, Akagi D, Soga Y, Iwakoshi S, Fujimura N, Kichikawa K. Clinical Impact of Stent Graft Thrombosis in Femoropopliteal Arterial Lesions. *JACC Cardiovasc Interv.* 2021;14:1137-1147.

29. Ogaki T, **Iida O**, Hata Y, Yamauchi N, Yokoi C, Takahara M, Terashi H, Mano T, Asada Y. The perioperative and long-term fates of patients with chronic limb-threatening ischaemia who underwent secondary major amputations. *Int Wound J*. 2021. doi: 10.1111/iwj.13594. Online ahead of print.
30. Kurata N, **Iida O**, Takahara M, Asai M, Masuda M, Okamoto S, Ishihara T, Nanto K, Mano T. Predictive Factors for Restenosis Following Stent-Supported Endovascular Therapy with Intravascular Ultrasound Evaluation for Femoropopliteal Chronic Total Occlusion. *J Vasc Interv Radiol*. 2021;32:712-720.e1.
31. Okuno S, Ishihara T, **Iida O**, Asai M, Masuda M, Okamoto S, Nanto K, Kanda T, Tsujimura T, Matsuda Y, Hata Y, Mano T. Two-Year Clinical Outcomes of Biodegradable Polymer vs. Durable Polymer Drug-Eluting Stent Implantation in Patients With End-Stage Renal Disease on Dialysis. *Circ Rep*. 2020;3:18-25.
32. Hata Y, **Iida O**, Inoue K, Mano T. Histological Evaluation of Self-Expandable Interwoven Stent After Implantation in the Femoropopliteal Artery. *Circ Rep*. 2020;2:766-767.
33. Toyoshima T, **Iida O**, Tsujimura T, Hata Y, Okamoto S, Mano T. A case of occluded femoropopliteal saphenous vein bypass with threatening limb ischemia treated with endovascular stent graft relining. *J Vasc Surg Cases Innov Tech*. 2020;7:74-77. eCollection 2021 Mar.
34. Takahara M, Soga Y, Fujihara M, Kawasaki D, Kozuki A, **Iida O**. Association of Age with Mortality Rate after Femoropopliteal Endovascular Therapy for Intermittent Claudication. *J Atheroscler Thromb*. 2021 Feb 27. doi: 10.5551/jat.62356. Online ahead of print.
35. Higashino N, **Iida O**, Hata Y, Asai M, Masuda M, Okamoto S, Ishihara T, Nanto K, Kanda T, Tsujimura T, Okuno S, Matsuda Y, Takahara M, Mano T. Impact of Longer Hemodialysis Vintage with Higher Serum Phosphorus Level on Clinical Outcomes in Patients with Chronic Limb-Threatening Ischemia Presenting Tissue Loss after Endovascular Therapy. *J Atheroscler Thromb*. 2021 Feb 14. doi: 10.5551/jat.60095. Online ahead of print.
36. Kadoya Y, Zen K, **Iida O**, Yamamoto Y, Kawasaki D, Yamauchi Y, Shintani Y, Sugano T, Yokoi H, Matoba S, Nakamura M; OMOTENASHI Investigators. Thrombotic Lesions are Associated with Poor Outcomes after Endovascular Treatment in Patients with Non-Acute Aortoiliac Total Occlusions. *J Atheroscler Thromb*. 2021 Feb 10. doi: 10.5551/jat.61150. Online ahead of print.
37. Matsuda Y, Masuda M, Asai M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Kanda T, Tsujimura T, Hata Y, Uematsu H, Mano T. Clinical Frailty Score Predicts Long-Term Mortality and Hospitalization Due to Heart Failure After Implantation of Cardiac Implantable Electric Device. *Circ J*. 2021;85:1341-1348.

38. Ishihara T, Inoue K, **Iida O**, Hata Y, Mano T. Pathological evaluation 18 years after bare-metal stent implantation in the superficial femoral artery. *J Cardiol Cases*. 2020;23:94-97.
39. Takahara M, **Iida O**, Soga Y, Azuma N, Nanto S; PRIORITY investigators. Clinical Impact of Measures for Frailty Severity in Poor-Risk Patients Undergoing Revascularization for Chronic Limb-Threatening Ischemia. *J Atheroscler Thromb*. 2021 Jan 30. doi: 10.5551/jat.61481. Online ahead of print.
40. Takahara M, **Iida O**, Soga Y, Kodama A, Terashi H, Azuma N; SPINACH study investigators. Current and Past Obesity in Japanese Patients with Critical Limb Ischemia Undergoing Revascularization. *J Atheroscler Thromb*. 2021;28:44-51.
41. Tomoi Y, Takahara M, Soga Y, Fujihara M, **Iida O**, Kawasaki D, Kozuki A, Ando K. Prognostic Value of the CHA<sub>2</sub>DS<sub>2</sub>-VASc Score after Endovascular Therapy for Femoral Popliteal Artery Lesions. *J Atheroscler Thromb*. 2021;28:1153-1160.
42. Tomoi Y, Soga Y, Okazaki J, **Iida O**, Shiraki T, Hiramori S, Ando K. Drug-coated stent implantation vs. bypass surgery for in-stent occlusion after femoropopliteal stenting. *Heart Vessels*. 2021;36:646-653.
43. Takahara M, **Iida O**, Soga Y, Kodama A, Terashi H, Suzuki K, Sugimoto I, Azuma N. Heterogeneity of Age and Its Associated Features in Patients with Critical Limb Ischemia. *Ann Vasc Dis*. 2020;13:300-307.
44. Yanaka K, Konishi A, Shinke T, Kozuki A, Kawamori H, Tsukiyama Y, **Iida O**, Kadotani M, Omori T, Hirata KI. Open-Label Multicenter Registry on the Outcomes of In-Stent Restenosis Treated by Balloon Angioplasty with Optical Frequency Domain Imaging in the Superficial Femoral Artery (ISLAND-SFA Study). *Ann Vasc Dis*. 2020;13:291-299.
45. Shiraki T, Takahara M, **Iida O**, Soga Y, Kodama A, Miyashita Y, Shintani Y, Endo M, Azuma N; SPINACH study investigators. Baseline and Updated Information on Nutritional Status in Patients With Chronic Limb Threatening Ischaemia Undergoing Revascularisation. *Eur J Vasc Endovasc Surg*. 2021;61:467-472.
46. Tsujimura T, Takahara M, **Iida O**, Soga Y, Katsuki T, Fujihara M, Kawasaki D, Kozuki A, Mano T. Clinical outcomes of polymer-free, paclitaxel-coated stents versus stent grafts in peripheral arterial disease patients with femoropopliteal artery lesions. *J Vasc Surg*. 2021;73:1998-2008.e1
47. Takahara M, Soga Y, Fujihara M, Kawasaki D, Kozuki A, **Iida O**. Inverse association of diabetes and dialysis with the severity of femoropopliteal lesions and chronic total occlusion: a cross-sectional study of 2056 cases. *BMC Cardiovasc Disord*. 2020;20:514.

48. Ishihara T, Sotomi Y, Tsujimura T, **Iida O**, Kobayashi T, Hamanaka Y, Omatsu T, Sakata Y, Higuchi Y, Mano T. Impact of diabetes mellitus on the early-phase arterial healing after drug-eluting stent implantation. *Cardiovasc Diabetol*. 2020;19:203.
49. Tsujimura T, Takahara M, **Iida O**, Kohsaka S, Soga Y, Fujihara M, Mano T, Ohya M, Shinke T, Amano T, Ikari Y; J-EVT and J-PCI investigators; Japanese Association of Cardiovascular Intervention and Therapeutics. In-Hospital Outcomes after Endovascular Therapy for Acute Limb Ischemia: A Report from a Japanese Nationwide Registry [J-EVT Registry]. *J Atheroscler Thromb*. 2021;28:1145-1152.
50. Müller-Hülsbeck S, Benko A, Soga Y, Fujihara M, **Iida O**, Babaev A, O'Connor D, Zeller T, Dulas DD, Diaz-Cartelle J, Gray WA. Two-Year Efficacy and Safety Results from the IMPERIAL Randomized Study of the Eluvia Polymer-Coated Drug-Eluting Stent and the Zilver PTX Polymer-free Drug-Coated Stent. *Cardiovasc Intervent Radiol*. 2021;44:368-375.
51. Shiraki T, **Iida O**, Okamoto S, Tsujimura T, Hata Y, Takahara M, Mano T, Sakata Y. Long-Term Clinical Outcomes after Endovascular Therapy for Anti-Centromere Antibody-Positive Patients with Critical Limb-Threatening Ischemia. *J Vasc Interv Radiol*. 2020;31:1993-1997.
52. Tsujimura T, Ishihara T, Okuno S, **Iida O**, Kurata N, Asai M, Masuda M, Okamoto S, Nanto K, Kanda T, Matsuda Y, Hata Y, Mano T. Angioscopic assessments and clinical outcomes one year after polymer-free biolimus A9-coated coronary stent implantation. *J Cardiol*. 2021;77:457-464.
53. Matsuda Y, Masuda M, Asai M, **Iida O**, Kanda T, Mano T. Bailout cryoballoon ablation for hemodynamically unstable atrial fibrillation in patients with decompensated heart failure. *HeartRhythm Case Rep*. 2020;6:685-689. eCollection 2020 Oct.
54. Dake MD, Fanelli F, Lottes AE, O'Leary EE, Reichert H, Jiang X, Fu W, **Iida O**, Zen K, Schermerhorn M, Zeller T, Ansel GM. Prediction Model for Freedom from TLR from a Multi-study Analysis of Long-Term Results with the Zilver PTX Drug-Eluting Peripheral Stent. *Cardiovasc Intervent Radiol*. 2021;44:196-206.
55. Ishihara T, Yamaji K, **Iida O**, Kohsaka S, Inohara T, Shinke T, Ando H, Amano T, Sakata Y, Mano T, Ikari Y. Impact of peripheral artery disease on short-term outcomes after percutaneous coronary intervention: A report from Japanese nationwide registry. *PLoS One*. 2020;15:e0240095. eCollection 2020.
56. Tsujimura T, Ishihara T, **Iida O**, Hata Y, Kurata N, Asai M, Masuda M, Okamoto S, Nanto K, Kanda T, Matsuda Y, Mano T. Angioscopic Assessments at 3 Months After Fluoropolymer-Based Paclitaxel-Eluting Stent Implantation for Femoropopliteal Endovascular Intervention. *Circ J*. 2020;84:1999-2005.
57. Matsuda Y, Masuda M, Asai M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Kanda T, Tsujimura T, Hata Y, Uematsu H, Mano T. A new clinical risk score for predicting the prevalence of low-

voltage areas in patients undergoing atrial fibrillation ablation. *J Cardiovasc Electrophysiol.* 2020;31:3150-3158.

58. Soga Y, **Iida O**, Urasawa K, Saito S, Jaff MR, Wang H, Ookubo H, Yokoi H. Three-Year Results of the IN.PACT SFA Japan Trial Comparing Drug-Coated Balloons With Percutaneous Transluminal Angioplasty. *J Endovasc Ther.* 2020;27:946-955.
59. Tsujimura T, Takahara M, **Iida O**, Yamauchi Y, Shintani Y, Sugano T, Yamamoto Y, Kawasaki D, Yokoi H, Miyamoto A, Mano T, Nakamura M. Intravascular Ultrasound Imaging During Aortoiliac Stenting: No Impact on Outcomes at 1 Year. *J Endovasc Ther.* 2021;28:139-145.
60. Yamauchi Y, Takahara M, Shintani Y, **Iida O**, Sugano T, Yamamoto Y, Kawasaki D, Fujihara M, Hirano K, Soga Y, Tsubakimoto Y, Yokoi H, Miyamoto A, Nakamura M. Three-Year Outcomes of Endovascular Therapy for Aortoiliac Lesions. *Circ Cardiovasc Interv.* 2020;13:e009168.
61. Hata Y, **Iida O**, Asai M, Masuda M, Okamoto S, Ishihara T, Nanto K, Kanda T, Tsujimura T, Okuno S, Matsuda Y, Takahara M, Mano T. Risk Stratification for 2-Year Mortality in Patients with Chronic Limb-Threatening Ischemia Undergoing Endovascular Therapy. *J Atheroscler Thromb.* 2021;28:477-482.
62. Ishihara T, **Iida O**, Takahara M, Tsujimura T, Okuno S, Kurata N, Asai M, Okamoto S, Nanto K, Mano T. Improved crossability with novel cutting balloon versus scoring balloon in the treatment of calcified lesion. *Cardiovasc Interv Ther.* 2021;36:198-207.
63. Tsuji A, Masuda M, Asai M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Kanda T, Tsujimura T, Matsuda Y, Okuno S, Hata Y, Mano T. Impact of the Temporal Relationship Between Atrial Fibrillation and Heart Failure on Prognosis After Ablation. *Circ J.* 2020;84:1467-1474.
64. Masuda M, Kanda T, Kurata N, Asai M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Tsujimura T, Matsuda Y, Hata Y, Uematsu H, Mano T. Clinical utility of local impedance monitoring during pulmonary vein isolation. *J Cardiovasc Electrophysiol.* 2020;31:2584-2591.
65. Tsukiyama Y, Konishi A, Shinke T, Kozuki A, Otake H, Kawamori H, Yanaka K, **Iida O**, Ishihara T, Inoue T, Iwasaki M, Kadotani M, Matsukawa N, Noutomi K, Kakei Y, Nanba I, Omori T, Shite J, Hirata KI. Open-label multicenter registry on the outcomes of peripheral arterial disease treated by balloon angioplasty with optical frequency domain imaging in superficial femoral artery and popliteal artery (OCEAN-SFA study). *Cardiovasc Interv Ther.* 2021;36:321-329.
66. Hata Y, **Iida O**, Takahara M, Asai M, Masuda M, Okamoto S, Ishihara T, Nanto K, Kanda T, Tsujimura T, Okuno S, Matsuda Y, Mano T. Infrapopliteal Anatomic Severity and Delayed Wound Healing in Patients With Chronic Limb-Threatening Ischemia in the Era of the Global Limb Anatomic Staging System. *J Endovasc Ther.* 2020;27:641-646.

67. Matsuda Y, Masuda M, Asai M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Kanda T, Tsujimura T, Okuno S, Hata Y, Uematsu H, Mano T. Comparison of Prognosis and Safety of Pacemaker Implantation in Patients Aged Less Than or 85 Years and Older. *J Interv Card Electrophysiol*. 2021;61:171-179.
68. Kurata N, Masuda M, Kanda T, Asai M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Tsujimura T, Matsuda Y, Hata Y, Mano T. Slow whole left atrial conduction velocity after pulmonary vein isolation predicts atrial fibrillation recurrence. *J Cardiovasc Electrophysiol*. 2020;31:1942-1949.
69. Utsunomiya M, Takahara M, **Iida O**, Soga Y, Hata Y, Shiraki T, Nagae A, Kato T, Kobayashi N, Suematsu N, Tasaki J, Horie K, Uchida D, Kodama A, Azuma N, Nakamura M. Limb-Based Patency After Surgical vs Endovascular Revascularization in Patients with Chronic Limb-Threatening Ischemia. *J Endovasc Ther*. 2020;27:584-594.
70. Okamoto S, **Iida O**, Takahara M, Hata Y, Asai M, Masuda M, Ishihara T, Nanto K, Kanda T, Tsujimura T, Okuno S, Matsuda Y, Mano T. A Global Vascular Guidelines-Based Bypass-Preferred Population and Their Surgical Risk in CLTI Patients Treated With Endovascular Therapy in a Real-World Practice. *J Endovasc Ther*. 2020;27:608-613.
71. Takahara M, **Iida O**, Kohsaka S, Soga Y, Fujihara M, Shinke T, Amano T, Ikari Y; J-EVT and J-PCI investigators. Presentation Pattern of Lower Extremity Endovascular Intervention versus Percutaneous Coronary Intervention. *J Atheroscler Thromb*. 2020;27:761-768.
72. Masuda M, Konishi S, Asai M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Kanda T, Tsujimura T, Matsuda Y, Okuno S, Tsuji A, Minamiguchi H, Mizuno H, Sakata Y, Mano T. Usefulness of an  $I_{Kr}$  blocker for ablation of non-pulmonary vein ectopies that are unmappable due to easily initiated atrial fibrillation. *J Interv Card Electrophysiol*. 2020;58:203-208.
73. Masuda M, Asai M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Kanda T, Tsujimura T, Matsuda Y, Okuno S, Hata Y, Mano T. Additional Low-Voltage-Area Ablation in Patients With Paroxysmal Atrial Fibrillation: Results of the Randomized Controlled VOLCANO Trial. *J Am Heart Assoc*. 2020;9:e015927.
74. Sawano M, Yamaji K, Kohsaka S, Inohara T, Numasawa Y, Ando H, **Iida O**, Shinke T, Ishii H, Amano T. Contemporary use and trends in percutaneous coronary intervention in Japan: an outline of the J-PCI registry. *Cardiovasc Interv Ther*. 2020;35:218-226.
75. Kanda T, Masuda M, Kurata N, Asai M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Tsujimura T, Okuno S, Matsuda Y, Hata Y, Mano T. A saline contrast-enhanced echocardiography-guided approach to cryoballoon ablation. *Pacing Clin Electrophysiol*. 2020;43:664-670.
76. Okuno S, **Iida O**, Inoue K, Asai M, Masuda M, Okamoto S, Ishihara T, Nanto K, Kanda T, Tsujimura T, Matsuda Y, Hata Y, Mano T. Very Late Intrastent Thrombotic Occlusion Caused



by Neoatherosclerosis After Bare-Metal Stent Implantation in the Superficial Femoral Artery: Insight From Histopathological Findings. *J Cardiol Cases*. 2020;21:172-175.

77. Yamauchi Y, Takahara M, **Iida O**, Shintani Y, Sugano T, Yamamoto Y, Kawasaki D, Fujihara M, Soga Y, Hirano K, Yokoi H, Miyamoto A, Nakamura M; OMOTENASHI investigators. Independent predictors of loss of primary patency at 1 year after aortoiliac stent implantation. *Heart Vessels*. 2020;35:614-619.
78. Suzuki K, **Iida O**, Yamauchi Y, Nakano M, Soga Y, Kawasaki D, Tazaki J, Yamaoka T, Suematsu N, Shintani Y, Miyashita Y, Inoue N, Meguro T. Impact of Diabetes Mellitus on Critical Limb Ischemia With Below the Knee Disease: Japan Below-the-Knee Artery Treatment Subanalysis. *Angiology*. 2020;71:444-451.
79. Fukunaga M, Fujii K, Takahara M, Fujihara M, Utsunomiya M, Sasaki S, Tsubakimoto Y, Shiraki T, Inoue T, Aihara H, Iwasaki Y, Azuma K, Nakata T, **Iida O**, Yokoi H, Yokoi Y, Kawasaki D. Assessment of Microvasculature After Intervention as a Predictor of Wound Healing in Patients With Tissue Loss. *JACC Cardiovasc Interv*. 2020;13:976-985.
80. Golzar J, Soga Y, Babaev A, **Iida O**, Kawasaki D, Bachinsky W, Park J, Prem JT, Vermassen F, Diaz-Cartelle J, Müller-Hülsbeck S, Gray WA. Effectiveness and Safety of a Paclitaxel-Eluting Stent for Superficial Femoral Artery Lesions up to 190 mm: One-Year Outcomes of the Single-Arm IMPERIAL Long Lesion Substudy of the Eluvia Drug-Eluting Stent. *J Endovasc Ther*. 2020;27:296-303.
81. Takahara M, **Iida O**, Soga Y, Kodama A, Terashi H, Utsunomiya M, Okazaki J, Azuma N, The Spinach Study Investigators. Duration from Wound Occurrence to Referral to a Vascular Center in Japanese Patients with Critical Limb Ischemia. *Ann Vasc Dis*. 2020;13:56-62.
82. Tomoi Y, Kuramitsu S, Soga Y, Fujihara M, **Iida O**, Kawasaki D, Ando K. Vascular Response After FP-PES Versus Polymer-Free Paclitaxel-Eluting Stent Implantation in Femoropopliteal Artery Lesions: A Serial Intravascular Ultrasound Study. *JACC Cardiovasc Interv*. 2020;13:535-537.
83. Kurata N, **Iida O**, Asai M, Masuda M, Okamoto S, Ishihara T, Nanto K, Mano T. Factors Influencing In-Stent Occlusion after Femoropopliteal Artery Stent Placement with Intravascular Ultrasound Evaluation. *J Vasc Interv Radiol*. 2020;31:213-220.
84. Soga Y, Fujihara M, **Iida O**, Kawasaki D, Hirano K, Yokoi H, Miyamoto A, Kichikawa K, Nakamura M, Ohki T, Diaz-Cartelle J, Gray WA, Müller-Hülsbeck S. Japanese Patients Treated in the IMPERIAL Randomized Trial Comparing Eluvia and Zilver PTX Stents. *Cardiovasc Intervent Radiol*. 2020;43:215-222.
85. Soga Y, Fujihara M, Tomoi Y, **Iida O**, Ishihara T, Kawasaki D, Ando K. One-Year Late Lumen Loss between A Polymer-Coated Paclitaxel-Eluting Stent (Eluvia) and a Polymer-Free

Paclitaxel-Coated Stent (Zilver PTX) for Femoropopliteal Disease. *J Atheroscler Thromb.* 2020;27:164-171.

86. Takahara M, Okuno S, Nakamura I, **Iida O**, Tsujimura T, Hata Y, Fujita Y, Haneda M. Prospective study on clinical characteristics of Japanese diabetic patients with chronic limb-threatening ischemia presenting Fontaine stage IV. *Diabetol Int.* 2019;11:33-40. eCollection 2020 Jan.
87. Dake MD, Ansel GM, Bosiers M, Holden A, **Iida O**, Jaff MR, Lottes AE, O'Leary EE, Saunders AT, Schermerhorn M, Yokoi H, Zeller T. Paclitaxel-Coated Zilver PTX Drug-Eluting Stent Treatment Does Not Result in Increased Long-Term All-Cause Mortality Compared to Uncoated Devices. *Cardiovasc Intervent Radiol.* 2020;43:8-19.
88. Azuma N, Takahara M, Kodama A, Soga Y, Terashi H, Tazaki J, Yamaoka T, Koya A, **Iida O**. Predictive Model for Mortality Risk Including the Wound, Ischemia, Foot Infection Classification in Patients Undergoing Revascularization for Critical Limb Ischemia. *Circ Cardiovasc Interv.* 2019;12:e008015.
89. Masuda M, Asai M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Kanda T, Tsujimura T, Matsuda Y, Okuno S, Hata Y, Mano T. Left atrial voltage mapping with a direction-independent grid catheter: comparison with conventional circular mapping catheter. *J Cardiovasc Electrophysiol.* 2019;30:2834-2840.
90. Takahara M, **Iida O**, Kohsaka S, Soga Y, Fujihara M, Shinke T, Amano T, Ikari Y; J-EVT and J-PCI investigators. Diabetes mellitus and other cardiovascular risk factors in lower-extremity peripheral artery disease versus coronary artery disease: an analysis of 1,121,359 cases from the nationwide databases. *Cardiovasc Diabetol.* 2019;18:155.
91. Tsujimura T, Ishihara T, **Iida O**, Asai M, Masuda M, Okamoto S, Nanto K, Kanda T, Okuno S, Matsuda Y, Mano T. Successful percutaneous retrieval of a detached microcatheter tip using the guide-extension catheter trapping technique: A case report. *J Cardiol Cases.* 2019;20:168-171. eCollection 2019 Nov.
92. Kanda T, Masuda M, Kurata N, Asai M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Tsujimura T, Okuno S, Matsuda Y, Hata Y, Mano T. Efficacy and safety of the cryoballoon-based atrial fibrillation ablation in patients aged  $\geq 80$  years. *J Cardiovasc Electrophysiol.* 2019;30:2242-2247.
93. Nanto K, **Iida O**, Fujihara M, Yokoi Y, Tomoi Y, Soga Y, Fujita M, Masuda M, Okamoto S, Ishihara T, Kanda T, Tsujimura T, Matsuda Y, Okuno S, Mano T. Five-Year Patency and its Predictors after Endovascular Therapy for Aortoiliac Occlusive Disease. *J Atheroscler Thromb.* 2019;26:989-996.

94. Shishehbor MH, Schneider PA, Zeller T, Razavi MK, Laird JR, Wang H, Tieché C, Parikh SA, **Iida O**, Jaff MR. Total IN.PACT drug-coated balloon initiative reporting pooled imaging and propensity-matched cohorts. *J Vasc Surg*. 2019;70:1177-1191.e9.
95. Katsuki T, Takahara M, Soga Y, Okamoto S, **Iida O**, Fujihara M, Kawasaki D, Ando K. Mortality Risk Following Application of a Paclitaxel-Coated Stent in Femoropopliteal Lesions. *J Endovasc Ther*. 2019;26:593-599.
96. Hata Y, **Iida O**, Okamoto S, Ishihara T, Nanto K, Tsujumura T, Takahara M, Mano T. Additional Risk Stratification Using Local and Systemic Factors for Patients with Critical Limb Ischaemia Undergoing Endovascular Therapy in the Wifi Era. *Eur J Vasc Endovasc Surg*. 2019;58:548-555.
97. Masuda M, Asai M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Kanda T, Tsujimura T, Matsuda Y, Okuno S, Tsuji A, Mano T. Comparison of ablation outcomes of the second ablation procedure for recurrent atrial fibrillation using an ultra-high-resolution mapping system and conventional mappings system. *Clin Cardiol*. 2019;42:966-973.
98. Tsukiyama Y, Shinke T, Ishihara T, Otake H, Terashita D, Kozuki A, Fukunaga M, Zen K, Horimatsu T, Fujii K, Shite J, Uematsu M, Takahara M, **Iida O**, Nanto S, Hirata KI. Vascular response to paclitaxel-eluting nitinol self-expanding stent in superficial femoral artery lesions: post-implantation angioscopic findings from the SHIMEJI trial (Suppression of vascular wall Healing after IMplantation of drug Eluting peripheral stent in Japanese patients with the Infra inguinal lesion: serial angioscopic observation). *Int J Cardiovasc Imaging*. 2019;35:1777-1784.
99. Tomoi Y, Soga Y, Takahara M, Fujihara M, **Iida O**, Kawasaki D, Ando K. Spot stenting versus full coverage stenting after endovascular therapy for femoropopliteal artery lesions. *J Vasc Surg*. 2019;70:1166-1176.
100. Fujii T, Ikari Y, Hashimoto H, Kadota K, Amano T, Uemura S, Takashima H, Nakamura M; J-PCI Investigators. (Kadota K, Shiode N, Tanaka N, Amano T, Uemura S, Akasaka T, Morino Y, Fujii K, Hikichi H, Amano T, Fujii K, Kohsaka S, Ishii H, Tanabe K, Ozaki Y, Sumitsuji S, **Iida O**, Hara H, Takashima H, Shirai S, Nanasato M, Inohara T, Ueda Y, Numasawa Y, Noma S; Collaborators) Post-interventional adverse event risk by vascular access site among patients with acute coronary syndrome in Japan: observational analysis with a national registry J-PCI database. *Cardiovasc Interv Ther*. 2019;34:297-304.
101. Tsujimura T, Takahara M, **Iida O**, Hiramori S, Hayakawa N, Karashima E, Miura T, Teramura M, Ichihashi K, Kojima T, Aihara H, Yamaoka T, Fujihara M, Tosaka A, Dojjiri T, Mano T, Soga Y. One-Year Clinical Outcomes following Implantation of Innova™ Self-Expanding Nitinol Stents in Patients with Peripheral Artery Diseases Presenting Femoropopliteal Artery Lesions. *J Atheroscler Thromb*. 2019;26:847-855.
102. Hata Y, **Iida O**, Mano T. Is Angiosome-Guided Endovascular Therapy Worthwhile? *Ann Vasc Dis*. 2019;12:315-318.

103. Soga Y, Takahara M, **Iida O**, Kodama A, Azuma N; SPINACH Investigators. Association between stress hormones and perioperative risk in patients with critical limb ischemia undergoing revascularization. *J Vasc Surg.* 2019;70:795-805.e1.
104. Takahara M, **Iida O**, Fujita Y, Haneda M. Clinical characteristics of Japanese diabetic patients with critical limb ischemia presenting Fontaine stage IV. *Diabetol Int.* 2018;10:231-235.
105. Ishihara T, Tsujimura T, Okuno S, **Iida O**, Asai M, Masuda M, Okamoto S, Nanto K, Kanda T, Matsuda Y, Mano T. Early- and middle-phase arterial repair following bioresorbable- and durable-polymer drug-eluting stent implantation: An angioscopic study. *Int J Cardiol.* 2019;285:27-31.
106. Matsuda Y, Masuda M, Asai M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Kanda T, Tsujimura T, Okuno S, Tsuji A, Mano T. A decapolar 25-mm-diameter circular mapping catheter with improved detection of pulmonary vein potential disappearance in cryoballoon ablation. *J Arrhythm.* 2019;35:535-541. eCollection 2019 Jun.
107. Masuda M, Asai M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Kanda T, Tsujimura T, Matsuda Y, Okuno S, Tsuji A, Mano T. Comparison of electrogram waveforms between a multielectrode mapping catheter and a linear ablation catheter. *Pacing Clin Electrophysiol.* 2019;42:515-520.
108. Okuno S, **Iida O**, Iida T, Takahara M, Yamaoka T, Kitano I, Asai M, Masuda M, Okamoto S, Ishihara T, Nanto K, Kanda T, Tsujimura T, Matsuda Y, Mano T. Comparison of Clinical Outcomes between Endovascular Therapy with Self-Expandable Nitinol Stent and Femoral-Popliteal Bypass for Trans-Atlantic Inter-Society Consensus II C and D Femoropopliteal Lesions. *Ann Vasc Surg.* 2019;57:137-143.
109. Matsuda Y, Masuda M, Asai M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Kanda T, Tsujimura T, Okuno S, Tsuji A, Mano T. Impact of Renal Dysfunction on Left Atrial Low-Voltage Areas in Patients With Atrial Fibrillation. *Circ J.* 2019;83:985-990.
110. Shiraki T, **Iida O**, Inoue K, Goto T, Mano T, Sakata Y. Very Late Histopathologic Evaluation of Neointimal Characteristics after Bare-Metal Stent Implantation in the Superficial Femoral Artery. *J Vasc Interv Radiol.* 2019;30:616-617.
111. Takahara M, **Iida O**, Soga Y, Kawasaki D, Fujihara M. Mortality risk of femoropopliteal paclitaxel-coated devices remains inconclusive. *Cardiovasc Interv Ther.* 2019;34:194-195.
112. Kurata N, Masuda M, Asai M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Kanda T, Tsujimura T, Matsuda Y, Okuno S, Ohashi T, Abe A, Mano T. Centrifugal wave-front propagation speed for localizing the atrial tachycardia origin. *Int J Cardiol.* 2019;279:96-99.
113. Kanda T, Masuda M, Kurata N, Matsuda Y, **Osamu I**, Asai M, Mano T. An unusual propagation pattern along the tricuspid annulus after cavo-tricuspid isthmus ablation: Insights

into posterior transverse conduction revealed by an ultra-high-resolution 3-dimensional mapping system. *J Cardiol Cases*. 2018;19:101-105. eCollection 2019 Mar.

114. Yamauchi Y, Takahara M, Shintani Y, **Iida O**, Sugano T, Yamamoto Y, Kawasaki D, Fujihara M, Hirano K, Yokoi H, Miyamoto A, Nakamura M; OMOTENASHI investigators. One-Year Outcomes of Endovascular Therapy for Aortoiliac Lesions. *Circ Cardiovasc Interv*. 2019 ;12:e007441.
115. Okuno S, Ishihara T, **Iida O**, Okamoto S, Nanto K, Mano T. Satisfactory arterial healing after second-generation drug-eluting stent implantation for segmental stenosis in a patient with Kawasaki disease. *Cardiovasc Interv Ther*. 2019;34:83-84.
116. Ishihara T, Awata M, **Iida O**, Fujita M, Masuda M, Okamoto S, Nanto K, Kanda T, Tsujimura T, Uematsu M, Mano T. Satisfactory arterial repair 1 year after ultrathin strut biodegradable polymer sirolimus-eluting stent implantation: an angioscopic observation. *Cardiovasc Interv Ther*. 2019;34:34-39.
117. Tokuda T, Hirano K, Soga Y, **Iida O**, Kawasaki D, Yamauchi Y, Suzuki K, Kamoi D, Tazaki J. Comparison of 3-year clinical outcomes after endovascular therapy for aortoiliac artery occlusive disease between patients with and without hemodialysis: Subanalysis of the REAL-AI registry. *Catheter Cardiovasc Interv*. 2018;92:1345-1351.
118. Kanda T, Uematsu M, Fujita M, **Iida O**, Masuda M, Okamoto S, Ishihara T, Nanto K, Tsujimura T, Matsuda Y, Okuno S, Mano T. A novel predictor of clinical outcomes in patients with heart failure with preserved left-ventricular ejection fraction: a pilot study. 2018 ;33:1490-1495.
119. Sunaga A, Masuda M, Asai M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Kanda T, Tsujimura T, Matsuda Y, Okuno S, Mano T. Pressure monitoring predicts pulmonary vein occlusion in cryoballoon ablation. *J Interv Card Electrophysiol*. 2018;53:115-121.
120. Suemitsu K, Shiraki T, **Iida O**, Kobayashi H, Matsuoka Y, Izumi M, Nakanishi T. Impact of Lesion Morphology on Durability After Angioplasty of Failed Arteriovenous Fistulas in Hemodialysis Patients. *J Endovasc Ther*. 2018;25:649-654.
121. Takahara M, **Iida O**, Soga Y, Azuma N, Nanto S; PRIORITY Investigators. Length and Cost of Hospital Stay in Poor-Risk Patients With Critical Limb Ischemia Undergoing Revascularization. *Circ J*. 2018;82:2634-2639.
122. Masuda M, Fujita M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Kanda T, Sunaga A, Tsujimura T, Matsuda Y, Ohashi T, Uematsu M. Pace-capture-guided ablation after contact-force-guided pulmonary vein isolation: results of the randomized controlled DRAGON trial. *Europace*. 2018;20:1451-1458.

123. Okuno S, Ishihara T, **Iida O**, Asai M, Masuda M, Okamoto S, Nanto K, Kanda T, Tsujimura T, Matsuda Y, Takahara M, Uematsu M, Mano T. Association of Subclinical Intrastent Thrombus Detected 9 Months After Implantation of 2nd-Generation Drug-Eluting Stent With Future Major Adverse Cardiac Events - A Coronary Angioscopic Study. *Circ J*. 2018;82:2299-2304.
124. Yokoi H, Ho M, Iwamoto S, Suzuki Y, Ansel GM, Azuma N, Handa N, **Iida O**, Ikeda K, Ikeno F, Ohura N, Rosenfield K, Rundback J, Terashi H, Uchida T, Yokoi Y, Nakamura M, Jaff MR. Design Strategies for Global Clinical Trials of Endovascular Devices for Critical Limb Ischemia (CLI) - A Joint USA-Japanese Perspective. *Circ J*. 2018;82:2233-2239.
125. Ishihara T, **Iida O**, Yutani C, Mano T. Near infrared spectroscopy, optical coherence tomography and angioscopy to unravel the mechanism of very late metallic drug-eluting stent thrombosis. *EuroIntervention*. 2018;14:686-687.
126. Soga Y, Hamasaki T, Edahiro R, **Iida O**, Inoue N, Suzuki K, Yokoi Y, Kawasaki D, Zen K, Urasawa K, Aodo K; STOP-IC investigators. Sustained Effectiveness of Cilostazol After Endovascular Treatment of Femoropopliteal Lesions: Midterm Follow-up From the Sufficient Treatment of Peripheral Intervention by Cilostazol (STOP-IC) Study. *J Endovasc Ther*. 2018;25:306-312.
127. Masuda M, Fujita M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Kanda T, Sunaga A, Tsujimura T, Matsuda Y, Ohashi T, Uematsu M. An E/e' ratio on echocardiography predicts the existence of left atrial low-voltage areas and poor outcomes after catheter ablation for atrial fibrillation. *Europace*. 2018;20:e60-e68.
128. Kurata N, **Iida O**, Shiraki T, Fujita M, Masuda M, Okamoto S, Ishihara T, Nanto K, Kanda T, Sunaga A, Tsujimura T, Takahara M, Mano T. Impact of Stent-to-Vessel Diameter Ratio on Restenosis in the Superficial Femoral Artery After Endovascular Therapy. *Circ J*. 2018;82:1412-1417.
129. Masuda M, Fujita M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Kanda T, Tsujimura T, Matsuda Y, Okuno S, Ohashi T, Tsuji A, Mano T. Left atrial low-voltage areas predict atrial fibrillation recurrence after catheter ablation in patients with paroxysmal atrial fibrillation. *Int J Cardiol*. 2018;257:97-101. doi: 10.1016/j.ijcard.2017.12.089.
130. Hata Y, **Iida O**, Asai M, Masuda M, Okamoto S, Ishihara T, Nanto K, Kanda T, Tsujimura T, Okuno S, Matsuda Y, Mano T. Endovascular entry closure using Amplatzer vascular plug for the aortic dissection in sub-acute phase. *J Cardiol Cases*. 2017;17:111-115. eCollection 2018 Apr.
131. Tsujimura T, **Iida O**, Fujita M, Masuda M, Okamoto S, Ishihara T, Nanto K, Kanda T, Okuno S, Matsuda Y, Fujihara M, Yokoi Y, Mano T. Two-year Clinical Outcomes Post Implantation of Epic™ Self-Expanding Nitinol Stents for the Aortoiliac Occlusive Disease in Patients with Peripheral Arterial Disease. *J Atheroscler Thromb*. 2018;25:344-349.



132. Ishihara T, **Iida O**, Fujita M, Masuda M, Okamoto S, Nanto K, Kanda T, Tsujimura T, Sunaga A, Awata M, Nanto S, Uematsu M. Comparison of early-phase arterial repair following cobalt-chrome everolimus-eluting stent and slow-release zotarolimus-eluting stent: an angioscopic study. *Cardiovasc Interv Ther*. 2018;33:163-168.
133. Masuda M, Asai M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Kanda T, Tsujimura T, Matsuda Y, Okuno S, Ohashi T, Tsuji A, Mano T. Centrifugal Wave Front Propagation Speed for Localizing the Origin of Ventricular Arrhythmias: Investigation Using a New Ultra-High-Resolution Mapping System. *JACC Clin Electrophysiol*. 2018;4:355-363.
134. Kanda T, Masuda M, Fujita M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Sunaga A, Tsujimura T, Matsuda Y, Ohashi T, Uematsu M. Comparison of the origin and coupling interval between ectopy with and without atrial fibrillation initiation. *J Cardiol*. 2018;71:59-64.
135. Tsuji Y, Shiraki T, **Iida O**, Tsuji Y, Kitano I, Sugimoto K, Okita Y. Impact of infrapopliteal endovascular treatment on the outcome of subsequent ipsilateral distal bypass for critical limb ischemia. *J Cardiovasc Surg (Torino)*. 2017;58:828-834.
136. Tsujimura T, **Iida O**, Ishihara T, Fujita M, Masuda M, Okamoto S, Nanto K, Kanda T, Sunaga A, Takahara M, Uematsu M. The impact of coronary artery disease and left ventricular ejection fraction on the prognosis of patients with peripheral artery disease. *Intern Med J*. 2017;47:1313-1316.
137. Masuda M, Fujita M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Kanda T, Tsujimura T, Matsuda Y, Okuno S, Ohashi T, Tsuji A, Mano T. The identification of conduction gaps after pulmonary vein isolation using a new electroanatomical mapping system. *Heart Rhythm*. 2017;14:1606-1614.
138. Kamioka N, Soga Y, Kuramitsu S, **Iida O**, Hirano K, Suzuki K, Kawasaki D, Yamaoka T, Suematsu N, Shintani Y, Miyashita Y, Takahashi H, Tsuchiya T, Shinozaki N, Okazaki S, Ando K. Clinical outcomes of balloon angioplasty alone versus nitinol stent implantation in patients with small femoropopliteal artery disease: Observations from the Retrospective Multicenter Analysis for Femoropopliteal Stenting (REAL-FP). *Catheter Cardiovasc Interv*. 2017;90:790-797.
139. Soga Y, Takahara M, **Iida O**, Yamauchi Y, Hirano K, Fukunaga M, Zen K, Suzuki K, Shintani Y, Miyashita Y, Tsuchiya T, Yamaoka T, Ando K. Efficacy of Cilostazol for Below-the-knee artery disease after Balloon Angioplasty in patients with severe limb ischemia (CABBAGE trial). *Ann Vasc Surg*. 2017;45:22-28.
140. Ishihara T, **Iida O**, Inoue K, Fujita M, Masuda M, Okamoto S, Nanto K, Kanda T, Tsujimura T, Sunaga A, Mano T, Uematsu M. Histological Evaluation of a Self-Expanding Stent-Graft 23 Months After Implantation in the Superficial Femoral Artery. *J Endovasc Ther*. 2017;24:746-750.

141. Tomoi Y, Soga Y, **Iida O**, Fujihara M, Ando K. Impact of Cilostazol Administration on Femoropopliteal In-Stent Restenosis. *J Endovasc Ther.* 2017;24:640-646.
142. Ishihara T, **Iida O**, Okamoto S, Fujita M, Masuda M, Nanto K, Shiraki T, Kanda T, Tsujimura T, Okuno S, Yanaka K, Uematsu M. Potential mechanisms of in-stent occlusion in the femoropopliteal artery: an angioscopic assessment. *Cardiovasc Interv Ther.* 2017;32:313-317.
143. Tsujimura T, Ishihara T, **Iida O**, Fujita M, Okamoto S, Uematsu M. A case complicated by floating fractured stent 141 months after nitinol stent implantation in the superficial femoral artery. *J Cardiol Cases.* 2017 Apr;16:38-40. eCollection 2017 Aug.
144. Masuda M, Fujita M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Kanda T, Sunaga A, Tsujimura T, Matsuda Y, Mano T. Association Between Local Bipolar Voltage and Conduction Gap Along the Left Atrial Linear Ablation Lesion in Patients With Atrial Fibrillation. *Am J Cardiol.* 2017;120:408-413.
145. Okamoto S, **Iida O**, Mano T. Current Perspective on Hemodialysis Patients with Peripheral Artery Disease. *Ann Vasc Dis.* 2017;10:88-91.
146. Fujihara M, Takahara M, Sasaki S, Nanto K, Utsunomiya M, **Iida O**, Yokoi Y. Angiographic Dissection Patterns and Patency Outcomes After Balloon Angioplasty for Superficial Femoral Artery Disease. *J Endovasc Ther.* 2017;24:367-375.
147. Sunaga A, Masuda M, Fujita M, **Iida O**, Kanda T, Matsuda Y, Morozumi T, Mano T, Uematsu M. Cardiac iodine-123-metaiodobenzylguanidine scintigraphy may be useful to identify pathologic from physiologic sinus bradycardia. *Pacing Clin Electrophysiol.* 2017;40:632-637.
148. Hiramori S, Soga Y, **Iida O**, Suzuki K, Hirano K, Kawasaki D, Shintani Y, Ando K. Relationship between clinical outcomes and vessel size in endovascular therapy for femoropopliteal lesions. *J Vasc Surg.* 2017;65:1690-1697.
149. Shiraki T, **Iida O**, Takahara M, Soga Y, Mii S, Okazaki J, Kuma S, Yamaoka T, Kamoi D, Shintani Y, Ishikawa T, Kitano I, Uematsu M. Comparison of Clinical Outcomes after Surgical and Endovascular Revascularization in Hemodialysis Patients with Critical Limb Ischemia. *J Atheroscler Thromb.* 2017;24:621-629.
150. Inohara T, Kohsaka S, Yamaji K, Amano T, Fujii K, Oda H, Uemura S, Kadota K, Miyata H, Nakamura M; J-PCI Registry Investigators. (Collaborators; Inohara T, Kohsaka S, Yamaji K, Amano T, Fujii K, Oda H, Uemura S, Kadota K, Miyata H, Nakamura M, Kadota K, Shiode N, Tanaka N, Amano T, Uemura S, Akasaka T, Morino Y, Fujii K, Hikichi H, Amano T, Fujii K, Kohsaka S, Ishii H, Tanabe K, Ozaki Y, Sumitsuji S, **Iida O**, Hara H, Takashima H, Shirai S, Nansato M, Inohara T, Ueda Y, Numasawa Y, Noma S.) Impact of Institutional and Operator Volume on Short-Term Outcomes of Percutaneous Coronary Intervention: A Report From the Japanese Nationwide Registry. *JACC Cardiovasc Interv.* 2017;10:918-927.

151. Masuda M, Fujita M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Kanda T, Sunaga A, Tsujimura T, Matsuda Y, Ohashi T, Uematsu M. Comparison of Left Atrial Voltage Between Sinus Rhythm and atrial Fibrillation in Association with Electrogram Wave Form. *Pacing Clin Electrophysiol.* 2017;40:559-567.
152. Zen K, Takahara M, **Iida O**, Soga Y, Kawasaki D, Nanto S, Yokoi H, Matoba S; ZEPHYR Investigators. Drug-eluting stenting for femoropopliteal lesions, followed by cilostazol treatment, reduces stent restenosis in patients with symptomatic peripheral artery disease. *J Vasc Surg.* 2017 65:720-725.
153. Utsunomiya M, Takahara M, **Iida O**, Yamauchi Y, Kawasaki D, Yokoi Y, Soga Y, Ohura N, Nakamura M; OLIVE Investigators. Wound Blush Obtainment Is the Most Important Angiographic Endpoint for Wound Healing. *JACC Cardiovasc Interv.* 2017;10:188-194.
154. Rundback JH, Armstrong EJ, Contos B, **Iida O**, Jacobs D, Jaff MR, Matsumoto AH, Mills JL Sr, Montero-Baker M, Pena C, Tallian A, Uematsu M, Wilkins LR, Shishehbor MH. Key Concepts in Critical Limb Ischemia: Selected Proceedings from the 2015 Vascular Interventional Advances Meeting. *Ann Vasc Surg.* 2017;38:191-205.
155. Suzuki K, Mizutani Y, Soga Y, **Iida O**, Kawasaki D, Yamauchi Y, Hirano K, Koshida R, Kamoi D, Tazaki J, Higashitani M, Shintani Y, Yamaoka T, Okazaki S, Suematsu N, Tsuchiya T, Miyashita Y, Shinozaki N, Takahashi H, Inoue N. Efficacy and Safety of Endovascular Therapy for Aortoiliac TASC D Lesions. *Angiology.* 2017;68:67-73.
156. Ishihara T, Takahara M, **Iida O**, Soga Y, Hirano K, Yamauchi Y, Zen K, Kawasaki D, Nanto S, Yokoi H, Uematsu M; ZEPHYR Investigators. Comparable 2-Year Restenosis Rates Following Subintimal and Intraluminal Drug-Eluting Stent Implantation for Femoropopliteal Chronic Total Occlusion. *J Endovasc Ther.* 2016;23:889-895.
157. Suzuki K, Takahara M, Shintani Y, Tanaka A, Soga Y, Yamaoka T, Tosaka A, Sasaki S, Kawasaki D, Tsuchiya T, Kozuki A, **Iida O**, Retrospective Multicenter Comparison of S.M.A.R.T. CONTROL and MISAGO Stents in Treatment of Femoropopliteal Lesions. *J Vasc Interv Radiol.* 2016;27:1642-1649.
158. Tomoi Y, Soga Y, Fujihara M, **Iida O**, Shintani Y, Zen K, Ando K. Outcomes of Endovascular Therapy for Upper Extremity Peripheral Artery Disease With Critical Hand Ischemia. Tomoi Y, Soga Y, Fujihara M, **Iida O**, Shintani Y, Zen K, Ando K. *J Endovasc Ther.* 2016;23:717-22.
159. Utsunomiya M, **Iida O**, Yamauchi Y, Nakano M, Soga Y, Kawasaki D, Takahara M, Nakamura M. Influence of Repeat Intervention on the Risk of Major Amputation After Infrapopliteal Angioplasty for Critical Limb Ischemia. *J Endovasc Ther.* 2016;23:710-6.
160. Okuno S, **Iida O**, Shiraki T, Fujita M, Masuda M, Okamoto S, Ishihara T, Nanto K, Kanda T, Takahara M, Uematsu M. Impact of Calcification on Clinical Outcomes After Endovascular

Therapy for Superficial Femoral Artery Disease: Assessment Using the Peripheral Artery Calcification Scoring System. *J Endovasc Ther.* 2016;23:731-7.

161. Masuda M, Fujita M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Kanda T, Shiraki T, Sunaga A, Matsuda Y, Uematsu M. Steerable versus non-steerable sheaths during pulmonary vein isolation: impact of left atrial enlargement on the catheter-tissue contact force. *J Interv Card Electrophysiol.* 2016;47:99-107.
162. Yanaka K, Ishihara T, **Iida O**, Okamoto S, Fujita M, Masuda M, Nanto K, Shiraki T, Kanda T, Uematsu M. Yellow Neointima Following Stent Implantation in the Superficial Femoral Artery on Angioscopy. *Circ J.* 2016;80:2249-51.
163. Ishihara T, **Iida O**, Shiraki T, Fujita M, Inoue K, Uematsu M. Arterial repair after bare-metal stent implantation in peripheral arteries is delayed compared to that in coronary arteries: A case report of pathological evaluation. *J Cardiol Cases.* 2016;14:74-77. eCollection 2016 Sep.
164. Suemitsu K, **Iida O**, Shiraki T, Suemitsu S, Murakami M, Miyamoto M, Izumi M, Nakanishi T. Predicting loss of patency after forearm loop arteriovenous graft. *J Vasc Surg.* 2016;64:395-401.
165. Murata N, Takahara M, Soga Y, Nakano M, Yamauchi Y, Zen K, Kawasaki D, Yokoi H, Tosaka A, Tanaka N, **Iida O**. Drug-Eluting Stent vs Percutaneous Transluminal Angioplasty for Treatment of Femoropopliteal In-Stent Restenosis: Results From a Retrospective 1-Year Multicenter Study. *J Endovasc Ther.* 2016;23:642-7.
166. Tsuchiya T, Takamura T, Soga Y, **Iida O**, Hirano K, Suzuki K, Yamaoka T, Miyashita Y, Kitayama M, Kajinami K. Clinical impact and risk stratification of balloon angioplasty for femoropopliteal disease in nitinol stenting era: Retrospective multicenter study using propensity score matching analysis. *SAGE Open Med.* 2016 Jul 18;4:2050312116660116.
167. Tomoi Y, Soga Y, **Iida O**, Shiraki T, Kobayashi Y, Hiramori S, Ando K. Impact of Drug-Eluting Stent Implantation for Femoropopliteal In-Stent Occlusion. *J Endovasc Ther.* 2016;23:461-7.
168. Shiraki T, **Iida O**, Suemitsu K, Tsuji Y, Uematsu M. Retrograde Approach Using Surgical Cutdown Technique for Limb Salvage in a Case of Critical Limb Ischemia With Severely Calcified Tibial Occlusive Disease. *Vasc Endovascular Surg.* 2016;50:295-8.
169. Okamoto S, **Iida O**, Takahara M, Yamauchi Y, Hirano K, Soga Y, Suzuki K, Uematsu M. Impact of Perioperative Complications After Endovascular Therapy in Diabetic Patients With Critical Limb Ischemia due to Isolated Infrapopliteal Lesions. *J Endovasc Ther.* 2016;23:371-7.

170. Masuda M, Fujita M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Kanda T, Shiraki T, Sunaga A, Matsuda Y, Uematsu M. Influence of underlying substrate on atrial tachyarrhythmias after pulmonary vein isolation. *Heart Rhythm*. 2016;13:870-8.
171. Hioki H, Miyashita Y, Shiraki T, **Iida O**, Uematsu M, Miura T, Ebisawa S, Ikeda U. Impact of deteriorated calcium-phosphate homeostasis on amputation-free survival after endovascular revascularization in patients with critical limb ischemia on hemodialysis. *Vasc Med*. 2016;21:137-43.
172. Soga Y, Takahara M, **Iida O**, Nakano M, Yamauchi Y, Zen K, Kawasaki D, Ando K. Propensity Score Analysis Comparing Clinical Outcomes of Drug-Eluting vs Bare Nitinol Stents in Femoropopliteal Lesions. *J Endovasc Ther*. 2016;23:33-9.
173. Higashimori A, **Iida O**, Yamauchi Y, Kawasaki D, Nakamura M, Soga Y, Zen K, Yokoi Y; OLIVE Investigators. Outcomes of One straight-line flow with and without pedal arch in patients with critical limb ischemia. *Catheter Cardiovasc Interv*. 2016;87:129-33.
174. Soga Y, Takahara M, **Iida O**, Suzuki K, Hirano K, Kawasaki D, Shintani Y, Yamaoka T, Ando K. Relationship Between Primary Patency and Lesion Length Following Bare Nitinol Stent Placement for Femoropopliteal Disease. *J Endovasc Ther*. 2015;22:862-7.
175. Kanda T, Masuda M, Sunaga A, Fujita M, **Iida O**, Okamoto S, Ishihara T, Watanabe T, Takahara M, Sakata Y, Uematsu M. Low left atrial appendage flow velocity predicts recurrence of atrial fibrillation after catheter ablation of persistent atrial fibrillation. *J Cardiol*. 2015;66:377-81.
176. Shiraki T, **Iida O**, Takahara M, Soga Y, Mii S, Okazaki J, Kuma S, Yamaoka T, Kamoi D, Shintani Y, Ishikawa T, Kitano I, Uematsu M. Predictors of 2-Year Mortality and Risk Stratification After Surgical or Endovascular Revascularization of Infrainguinal Artery Disease in Hemodialysis Patients With Critical Limb Ischemia. *J Endovasc Ther*. 2015;22:719-24.
177. Shiraki T, **Iida O**, Okamoto S, Ishihara T, Fujita M, Uematsu M. Usefulness of Endovascular Therapy for Penile Gangrene Secondary to Calciphylaxis: A Case Report. *Ann Vasc Surg*. 2015;29:1451.e11-5.
178. Okamoto S, **Iida O**, Nakamura M, Yamauchi Y, Fukunaga M, Yokoi Y, Soga Y, Zen K, Hirano K, Suematsu N, Suzuki K, Shintani Y, Miyashita Y, Urasawa K, Kitano I, Yamaoka T, Ohura N, Hamasaki T, Uematsu M, Nanto S; on behalf of the OLIVE Investigators. Postprocedural Skin Perfusion Pressure Correlates With Clinical Outcomes 1 Year After Endovascular Therapy for Patients With Critical Limb Ischemia. *Angiology*. 2015;66:862-6.
179. Shintani Y, Soga Y, Takahara M, **Iida O**, Kawasaki D, Yamauchi Y, Suzuki K, Hirano K, Kawasaki T. Clinical Outcomes of SMART Versus Luminexx Nitinol Stent Implantation for Aortoiliac Artery Disease: A Propensity Score-Matched Multicenter Study. *Angiology*. 2015;66:875-81.

180. Sunaga A, Masuda M, Kanda T, Fujita M, **Iida O**, Okamoto S, Ishihara T, Matsuda Y, Watanabe T, Sakata Y, Uematsu M. A low fibrillatory wave amplitude predicts sinus node dysfunction after catheter ablation in patients with persistent atrial fibrillation. *J Interv Card Electrophysiol.* 2015;43:253-61.
181. Nanto K, **Iida O**, Takahara M, Soga Y, Suzuki K, Hirano K, Kawasaki D, Shintani Y, Suematsu N, Yamaoka T, Uematsu M. Effect of Cilostazol Following Endovascular Intervention for Peripheral Artery Disease. *Angiology.* 2015;66:774-8.
182. Nakano M, Hirano K, **Iida O**, Yamauchi Y, Soga Y, Kawasaki D, Tazaki J, Suzuki K, Fujiwara M, Yamaoka T. Clinical efficacy of infrapopliteal endovascular procedures for hemodialysis patients with critical limb ischemia. *Ann Vasc Surg.* 2015;29:1225-34.
183. Nakano M, Hirano K, Yamauchi Y, **Iida O**, Soga Y, Kawasaki D, Yamaoka T, Suematsu N, Suzuki K. Three-year clinical outcome after infrapopliteal angioplasty for critical limb ischemia in hemodialysis patients with minor or major tissue loss. *Catheter Cardiovasc Interv.* 2015;86:289-98.
184. Takahara M, **Iida O**, Soga Y, Kodama A, Azuma N; SPINACH study investigators. Absence of Preceding Intermittent Claudication and its Associated Clinical Features in Patients with Critical Limb Ischemia. *J Atheroscler Thromb.* 2015;22:718-25.
185. Tsuchiya T, **Iida O**, Shiraki T, Soga Y, Hirano K, Suzuki K, Yamaoka T, Miyashita Y, Kitayama M, Kajinami K. Clinical characteristics of patients with Rutherford category IV, compared with V and VI. *SAGE Open Med.* 2015;3:2050312115597087.
186. Kanda T, Masuda M, Sunaga A, Fujita M, **Iida O**, Okamoto S, Ishihara T, Nanto K, Shiraki T, Sera F, Uematsu M. Fabry cardiomyopathy presenting with a high defibrillation threshold: A short case report. *J Arrhythm.* 2015;31:170-1.
187. Kobayashi Y, Sakaki M, Yasuoka T, **Iida O**, Dohi T, Uematsu M. Endovascular repair with contralateral external-to-internal iliac artery bypass grafting: a case series. *BMC Res Notes.* 2015;8:183.
188. Sato K, **Iida O**, Takahara M, Soga Y, Suzuki K, Tanigawa T, Ito M, Uematsu M. Effect of perioperative complications after endovascular therapy in patients with peripheral artery disease due to femoropopliteal lesions. *J Vasc Surg.* 2015;61:1272-7.
189. Soga Y, Takahara M, **Iida O**, Mii S, Okazaki J, Nakano M, Yamauchi Y, Ando K. Bypass Surgery vs. Drug-Eluting Stent for Trans-Atlantic Inter-Society Consensus-II (TASCII) C or D Femoropopliteal Lesions. *Circ J.* 2015;79:2688-95.
190. Kanda T, Fujita M, **Iida O**, Masuda M, Okamoto S, Ishihara T, Nanto K, Shiraki T, Takahara M, Sakata Y, Uematsu M. Novel Echocardiographic Approach to the Accurate Measurement



of Pulmonary Vascular Resistance Based on a Theoretical Formula in Patients With Left Heart Failure - Pilot Study. *Circ J.* 2015;79:2408-13.

191. Ishihara T, Fujita M, **Iida O**, Tsujimura T, Uematsu M. Rupture of Calcified Nodule 105 Months After Sirolimus-Eluting Stent Implantation Observed on Coronary Angioscopy and Optical Frequency Domain Imaging In Vivo. *Circ J.* 2015;79:2278-9.
192. Kawasaki D, **Iida O**, Fukunaga M, Kato M, Ohkubo N. Wire Passages of 0.035-inch Looped Wire Technique for Femoropopliteal Long Total Occlusions. *J Atheroscler Thromb.* 2015;22:1071-9.
193. Soga Y, **Iida O**, Takahaera M, Hirano K, Suzuki K, Kawasaki D. Beta-blocker Treatment Does Not Worsen Critical Limb Ischemia in Patients Receiving Endovascular Therapy. *J Atheroscler Thromb.* 2015;22:481-9.
194. Suematsu N, **Iida O**, Takahara M, Yamauchi Y, Soga Y, Nakano M, Hirano K, Kawasaki D, Yamaoka T, Suzuki K, Shintani Y, Miyashita Y, Tazaki J, Meno H, Inou T. Prognostic Factors in Hemodialysis Patients Undergoing Endovascular Treatment for Critical Limb Ischemia due to Isolated Below-the-Knee Disease. *J Atheroscler Thromb.* 2015;22:404-14.
195. Ishihara T, **Iida O**, Awata M, Nanto K, Shiraki T, Okamoto S, Iida T, Fujita M, Watanabe T, Nanto S, Uematsu M. Extensive arterial repair one year after paclitaxel-coated nitinol drug-eluting stent vs. bare-metal stent implantation in the superficial femoral artery. *Cardiovasc Interv Ther.* 2015;30:51-6.
196. Shiraki T, **Iida O**, Takahara M, Soga Y, Yamauchi Y, Hirano K, Kawasaki D, Fujihara M, Utsunomiya M, Tazaki J, Yamaoka T, Shintani Y, Suematsu N, Suzuki K, Miyashita Y, Tsuchiya T, Uematsu M. Predictors of Delayed Wound Healing after Endovascular Therapy of Isolated Infrapopliteal Lesions Underlying Critical Limb Ischemia in Patients with High Prevalence of Diabetes Mellitus and Hemodialysis. *Eur J Vasc Endovasc Surg.* 2015;49:565-73.
197. Murata N, Soga Y, **Iida O**, Yamauchi Y, Hirano K, Kawasaki D, Fujihara M, Tomoi Y. Complex relationship of body mass index with mortality in patients with critical limb ischemia undergoing endovascular treatment. *Eur J Vasc Endovasc Surg.* 2015;49:297-305.
198. Soga Y, **Iida O**, Takahara M, Hirano K, Suzuki K, Kawasaki D, Miyashita Y, Tsuchiya T. Two-year life expectancy in patients with critical limb ischemia. *JACC Cardiovasc Interv.* 2014;7:1444-9.
199. Azuma N, **Iida O**, Takahara M, Soga Y, Kodama A. Surgical reconstruction versus peripheral intervention in patients with critical limb ischemia - a prospective multicenter registry in Japan: The SPINACH study design and rationale. *Vascular.* 2014;22:411-20.

200. Iida T, **Iida O**, Okamoto S, Dohi T, Nanto K, Uematsu M. Endovascular therapy with novel high anterior tibial artery puncture for limb salvage in a case of critical lower limb ischemia. *Cardiovasc Interv Ther.* 2014;29:363-7.
201. Shiraki T, **Iida O**, Takahara M, Okamoto S, Kitano I, Tsuji Y, Terashi H, Uematsu M. Predictive scoring model of mortality after surgical or endovascular revascularization in patients with critical limb ischemia. *J Vasc Surg.* 2014;60:383-9.
202. Takahara M, Kaneto H, Katakami N, **Iida O**, Matsuoka TA, Shimomura I. Effect of sarpogrelate treatment on the prognosis after endovascular therapy for critical limb ischemia. *Heart Vessels.* 2014;29:563-7.
203. Miura T, Soga Y, Miyashita Y, **Iida O**, Kawasaki D, Hirano K, Suzuki K, Ikeda U. Five-year prognosis after endovascular therapy in claudicant patients with iliofemoral artery disease. *J Endovasc Ther.* 2014;21:381-8.
204. Soga Y, Mii S, **Iida O**, Okazaki J, Kuma S, Hirano K, Suzuki K, Kawasaki D, Yamaoka T, Kamoi D, Shintani Y. Propensity Score Analysis of Clinical Outcome After Bypass Surgery vs. Endovascular Therapy for Infrainguinal Artery Disease in Patients With Critical Limb Ischemia. *J Endovasc Ther.* 2014;21:243-53.
205. Dohi T, **Iida O**, Soga Y, Hirano K, Suzuki K, Takahara M, Uematsu M, Nanto S. Incidence, predictors, and prognosis of in-stent occlusion after endovascular treatment with nitinol stents for femoropopliteal lesions. *J Vasc Surg.* 2014;59:1009-1015.e1.
206. Aihara H, Soga Y, **Iida O**, Suzuki K, Tazaki J, Shintani Y, Miyashita Y; REAL-AI Registry Investigators. Long-Term Outcomes of Endovascular Therapy for Aortoiliac Bifurcation Lesions in the Real-AI Registry. *J Endovasc Ther.* 2014;21:25-33.
207. Soga Y, **Iida O**, Suzuki K, Hirano K, Kawasaki D, Shintani Y, Suematsu N, Yamaoka T, Tsuchiya T, Miyashita Y, Okazaki S, Shinozaki N, Takahashi H. Clinical Impact of Bisoprolol Versus Carvedilol in Patients Undergoing Femoropopliteal Stenting. *J Atheroscler Thromb.* 2014;21:691-702.
208. Ishihara T, Awata M, Fujita M, Watanabe T, **Iida O**, Ishida Y, Nanto S, Uematsu M. Angioscopic Assessment of Peri-Stent Contrast Staining Following Drug-Eluting Stent Implantation. *Circ J.* 2013;78:122-7.
209. Ishihara T, **Iida O**, Awata M, Shiraki T, Nanto S, Uematsu M. Angioscopically apparent large thrombus and uncovered stent struts 6 months after late stent thrombosis of a paclitaxel-coated nitinol drug-eluting stent implanted in the superficial femoral artery. *Cardiovasc Interv Ther.* 2014;29:82-5.

210. Watanabe T, Fujita M, Awata M, **Iida O**, Okamoto S, Ishihara T, Uematsu M. Integrity of stent polymer layer after drug-eluting stent implantation: in vivo comparison of sirolimus-, paclitaxel-, zotarolimus- and everolimus-eluting stents. *Cardiovasc Interv Ther.* 2014;29:4-10.
211. Soga Y, **Iida O**, Suzuki K, Hirano K, Kawasaki D, Shintani Y, Suematsu N, Yamaoka T. Initial and 3-year results after subintimal versus intraluminal approach for long femoropopliteal occlusion treated with a self-expandable nitinol stent. *J Vasc Surg.* 2013;58:1547-55.
212. Mizoguchi H, **Iida O**, Dohi T, Tomoda K, Kimura H, Inoue K, Iwata T, Tei K, Miura T. Abdominal Aortic Aneurysmal and Endovascular Device Infection With Iliopsoas Abscess Caused by Mycobacterium Bovis as a Complication of Intravesical Bacillus Calmette-Guérin Therapy. *Ann Vasc Surg.* 2013;27:1186.e1-5.
213. Tomoi Y, Soga Y, **Iida O**, Hirano K, Suzuki K, Kawasaki D, Yamauchi Y, Miyashita Y, Tazaki J, Nobuyoshi M. Efficacy of statin treatment after endovascular therapy for isolated below-the-knee disease in patients with critical limb ischemia. *Cardiovasc Interv Ther.* 2013;28:374-82.
214. Dohi T, **Iida O**, Okamoto S, Nanto K, Nanto S, Uematsu M. Mid-term clinical outcome following endovascular therapy in patients with chronic aortic occlusion. *Cardiovasc Interv Ther.* 2013;28:327-32.
215. Sakamoto Y, Hirano K, **Iida O**, Soga Y, Suzuki K, Muramatsu T, Tsukahara R. Five-year outcomes of self-expanding nitinol stent implantation for chronic total occlusion of the superficial femoral and proximal popliteal artery. *Catheter Cardiovasc Interv.* 2013;82:E251-6.
216. Ishihara T, **Iida O**, Tosaka A, Soga Y, Sakamoto Y, Hirano K, Nanto S, Uematsu M. Severity of Coronary Artery Disease Affects Prognosis of Patients With Peripheral Artery Disease. *Angiology.* 2013;64:417-22.
217. Soga Y, Tomoi Y, Sato K, **Iida O**, Yokoi H. Clinical outcome after endovascular treatment for isolated common femoral and popliteal artery disease. *Cardiovasc Interv Ther.* 2013;28:250-7.
218. Ishihara T, Awata M, Fujita M, Watanabe T, **Iida O**, Ishida Y, Nanto S, Uematsu M. Very late stent thrombosis 5 years after implantation of a sirolimus-eluting stent observed by angioscopy and optical coherence tomography. *JACC Cardiovasc Interv.* 2013;6:e28-30.
219. Nakano M, Hirano K, **Iida O**, Soga Y, Kawasaki D, Suzuki K, Miyashita Y. Prognosis of Critical Limb Ischemia in Hemodialysis Patients After Isolated Infrapopliteal Balloon Angioplasty: Results From the Japan Below-the-Knee Artery Treatment (J-BEAT) Registry. *J Endovasc Ther.* 2013;20:113-24.
220. Takahara M, **Iida O**, Soga Y, Hirano K, Yamaoka T, Kawasaki D, Suzuki K, Suematsu N, Shintani Y, Miyashita Y, Kaneto H, Shimomura I. Seasonal Variation in Critical Limb Ischemia Requiring Endovascular Therapy: An Analysis of a Multicenter Database of Japanese Patients

with Critical Limb Ischemia Undergoing Endovascular Therapy. *J Atheroscler Thromb.* 2013;20:726-32.

221. Ishihara T, **Iida O**, Awata M, Nanto K, Nanto S, Uematsu M. Angioscopic Assessment of Early Phase Arterial Repair After Paclitaxel-Coated Nitinol Drug-Eluting Stent Implantation in the Superficial Femoral Artery. *Circ J.* 2013;77:1838-43.
222. Ishihara T, **Iida O**, Awata M, Fujita M, Watanabe T, Ishida Y, Nanto S, Uematsu M. Two-Month Angioscopic Evaluation of Superficial Femoral Artery Treated With Paclitaxel-Coated Nitinol Drug-Eluting Stent. *Circ J.* 2013;77:1880-2.
223. Ishihara T, Awata M, Sera F, Fujita M, Watanabe T, **Iida O**, Ishida Y, Nanto S, Uematsu M. Arterial Repair 4 Months After Zotarolimus-Eluting Stent Implantation Observed on Angioscopy. *Circ J.* 2013;77:1186-92.
224. Soga Y, **Iida O**, Kawasaki D, Hirano K, Yamaoka T, Suzuki K. Impact of Cilostazol on Angiographic Restenosis after Balloon Angioplasty for Infrapopliteal Artery Disease in Patients with Critical Limb Ischemia. *Eur J Vasc Endovasc Surg.* 2012;44:577-81.
225. Takahara M, Kaneto H, **Iida O**, Katakami N, Matsuoka TA, Ikeda M, Shimomura I. Association of diabetes and hemodialysis with ankle pressure and ankle-brachial index in Japanese patients with critical limb ischemia. *Diabetes Care.* 2012;35:2000-4.
226. Ishihara T, **Iida O**, Okamoto S, Dohi T, Sato K, Nanto K, Fujita M, Watanabe T, Awata M, Sera F, Tanaka N, Ishida Y, Nanto S, Uematsu M. Successful outside-the-stent stenting for in-stent chronic total occlusion in the common iliac artery. *Cardiovasc Interv Ther.* 2012;27:131-6.
227. Soga Y, **Iida O**, Hirano K, Suzuki K, Yokoi H, Nobuyoshi M. Restenosis after stent implantation for superficial femoral artery disease in patients treated with cilostazol. *Catheter Cardiovasc Interv.* 2012;79:541-8.
228. Tosaka A, Soga Y, **Iida O**, Ishihara T, Hirano K, Suzuki K, Yokoi H, Nanto S, Nobuyoshi M. Classification and clinical impact of restenosis after femoropopliteal stenting. *J Am Coll Cardiol.* 2012;59:16-23.
229. Soga Y, **Iida O**, Kawasaki D, Yamauchi Y, Suzuki K, Hirano K, Koshida R, Kamoi D, Tazaki J, Higashitani M, Shintani Y, Yamaoka T, Okazaki S, Suematsu N, Tsuchiya T, Miyashita Y, Shinozaki N, Takahashi H; on behalf of REAL-AI investigators. Contemporary Outcomes After Endovascular Treatment for Aorto-Iliac Artery Disease. *Circ J.* 2012;76:2697-704.
230. Soga Y, **Iida O**, Hirano K, Suzuki K, Kawasaki D, Miyashita Y, Tsuchiya T, Nobuyoshi M. Impact of cilostazol after endovascular treatment for infrainguinal disease in patients with critical limb ischemia. *J Vasc Surg.* 2011;54:1659-67.

231. Soga Y, **Iida O**, Hirano K, Suzuki K, Tosaka A, Yokoi H, Nobuyoshi M. Utility of new classification based on clinical and lesional factors after self-expandable nitinol stenting in the superficial femoral artery. *J Vasc Surg.* 2011;54:1058-66.
232. Tsuji Y, Kitano I, **Iida O**, Kajita S, Sawada K, Nanto S. Popliteal pseudoaneurysm caused by stent fracture. *Ann Vasc Surg.* 2011;25:840.e5-8.
233. Bito Y, Sakaki M, **Iida O**, Inoue K, Yoshioka Y, Mizoguchi H. Clinical management of lower limb ischemia secondary to a persistent sciatic artery aneurysm: report of a case. *Surg Today.* 2011;41:402-5.
234. Takahara M, Kaneto H, **Iida O**, Katakami N, Sakamoto F, Matsuoka TA, Ikeda M, Shimomura I. No association of diabetic duration or insulin use with the prognosis of critical limb ischemia after endovascular therapy. *J Atheroscler Thromb.* 2011;18:1102-9.
235. Awata M, Uematsu M, Sera F, Ishihara T, Watanabe T, Fujita M, Onishi T, **Iida O**, Ishida Y, Nanto S, Nagata S. Angioscopic assessment of arterial repair following biodegradable polymer-coated biolimus A9-eluting stent implantation. - Comparison with durable polymer-coated sirolimus-eluting stent-. *Circ J.* 2011;75:1113-9.
236. Suzuki K, **Iida O**, Soga Y, Hirano K, Inoue N, Uematsu M, Yokoi H, Muramatsu T, Nanto S, Nobuyoshi M, Meguro T. Long-term results of the S.M.A.R.T. Control(TM) stent for superficial femoral artery lesions, J-SMART registry. *Circ J.* 2011;75:939-44.
237. Miyashita Y, Saito S, Miyamoto A, **Iida O**, Nanto S. Cilostazol increases skin perfusion pressure in severely ischemic limbs. *Angiology.* 2011;62:15-7.
238. Takahara M, Kaneto H, **Iida O**, Gorogawa S, Katakami N, Matsuoka TA, Ikeda M, Shimomura I. The influence of glycemic control on the prognosis of Japanese patients undergoing percutaneous transluminal angioplasty for critical limb ischemia. *Diabetes Care.* 2010;33:2538-42.
239. Terashi H, Iwayama T, **Iida O**, Kitano I, Tsuji Y. Dynamic skin perfusion pressure: a new measure of assessment for wound healing capacity and alternative angiosome in critical limb ischemia. *Plast Reconstr Surg.* 2010;126:215e-218e.
240. Onishi T, Uematsu M, Watanabe T, Fujita M, Awata M, **Iida O**, Sera F, Hirano Y, Nanto S, Nagata S. Objective interpretation of dobutamine stress echocardiography by diastolic dyssynchrony imaging: a practical approach. *J Am Soc Echocardiogr.* 2010 ;23:1103-8.
241. Soga Y, **Iida O**, Hirano K, Yokoi H, Nanto S, Nobuyoshi M. Mid-term clinical outcome and predictors of vessel patency after femoropopliteal stenting with self-expandable nitinol stent. *J Vasc Surg.* 2010;52:608-15.

242. Awata M, Nanto S, Uematsu M, Morozumi T, Watanabe T, Onishi T, **Iida O**, Sera F, Minamiguchi H, Kotani J, Nagata S. Heterogeneous arterial healing in patients following paclitaxel-eluting stent implantation: comparison with sirolimus-eluting stents. *JACC Cardiovasc Interv.* 2009;2:453-8.
243. Onishi T, Uematsu M, Nanto S, Morozumi T, Watanabe T, Awata M, **Iida O**, Sera F, Nagata S. Detection of diastolic abnormality by dyssynchrony imaging: correlation with coronary artery disease in patients presenting with visibly normal wall motion. *Circ J.* 2009;73:125-31.
244. Awata M, Nanto S, Uematsu M, Morozumi T, Watanabe T, Onishi T, **Iida O**, Sera F, Kotani J, Horii M, Nagata S. Angioscopic comparison of neointimal coverage between zotarolimus- and sirolimus-eluting stents. *J Am Coll Cardiol.* 2008;52:789-90.
245. Awata M, Kotani J, Uematsu M, Morozumi T, Watanabe T, Onishi T, **Iida O**, Sera F, Nanto S, Horii M, Nagata S. Serial angioscopic evidence of incomplete neointimal coverage after sirolimus-eluting stent implantation: comparison with bare-metal stents. *Circulation.* 2007;116:910-6.
246. Onishi T, Uematsu M, Nanto S, **Iida O**, Morozumi T, Kotani J, Awata M, Nagata S. Positive isovolumic relaxation velocity detected by a spectral tissue Doppler mapping technique as an indicator of coronary artery disease: a prospective study. *J Am Soc Echocardiogr.* 2007;20:158-64.
247. Takahara M, Kaneto H, **Iida O**, Gorogawa S, Ikeda M. High prevalence of glucose intolerance in Japanese patients with peripheral arterial disease. *Diabetes Res Clin Pract.* 2011;91:e24-5.



## Abstract

---

1. Long-term Clinical Efficacy of Endovascular Therapy for Patients with Critical Limb Ischemia due to Pure Isolated Infrapopliteal Lesions. *Circulation* 2012; (Abstract 13627)
2. Angiographic Restenosis and Its Clinical Impact after Below-the-knee Angioplasty. *J Am Coll Cardiol* 2012; 61th ACC (Abstract E2087)
3. Long-term Clinical Outcomes after Endovascular Therapy for Patients with Critical Limb Ischemia due to Isolated Below the Knee Lesions. *Circulation* 2011; (Abstract 11141)
4. Impact of Angiosome in Endovascular Therapy (EVT) on the Limb Salvage for the Patients with Critical Limb Ischemia (CLI) Presenting with Isolated Infrapopliteal Lesions. *J Am Coll Cardiol* 2011; 60th ACC (Abstract E1667)
5. Time Course of the Restenosis Following Nitinol Stenting in the Superficial Femoral Artery and the Factors Associated with Early and Late Restenoses. *Circulation* 2010 (Abstract 12185)
6. Stent Fracture, TASC II CD lesion as Restenosis Factors, and Cilostazol as a Negative-restenosis Factor Within a Year Following Nitinol Stent Implantation in the Superficial Femoral Artery. *J Am Coll Cardiol* 2010; 59th ACC (Abstract .E1688)
7. Acquiring Direct Flow Based on the Angiosome Concept in Endovascular Therapy Saves More Limbs than Increasing the Number of Patent Run-off Vessels in Patients with Critical Limb Ischemia. *Circulation* 2009 (Abstract 4427)
8. Factors Differ Between Early and Late Restenosis Following Nitinol Stent Implantation in the Superficial Femoral Artery. *Circulation* 2009 (Abstract 4424)
9. Stent Fracture does not Affect Long Term Patency in the Femoro-popliteal Artery: Four Years Experience. *Circulation* 2008 (Abstract 6189)
10. Low Ejection Fraction and Poor Performance Status Negate the Beneficial Effects of Complete Revascularization in Patients with Critical Limb Ischemia. *J Am Coll Cardiol* 2008; 57th (Abstract)
11. Complete Revascularization to the Pedal Arch may Improve the Survival of Patient with Critical Limb Ischemia. *Circulation* 2007 (Abstract 3228)
12. Cilostazol Attenuates Angiographic Restenosis in Patients with Femoro-popliteal Lesions *Circulation* 2007 (Abstract 3224)
13. Morphology of the Stent Fracture Influences the Outcome Following Superficial Femoral Artery Stenting. *J Am Coll Cardiol* 2007; 56th (Abstract)

14. Successful Revascularization of at Least a Branch of the Tibial Artery to the Pedal Arch Contributes Limb Salvage in Critical Limb Ischemia. *J Am Coll Cardiol* 2007; 56th (Abstract)
15. Cilostazol Reduces Target Lesion Revascularization in the Femoro-Popliteal Lesions in Patients Complaining of Claudication. *Circulation* 2005 (Abstract 2242)
16. Exercise Adversely Affects Stent Fracture in the Superficial Femoral Artery. *Circulation* 2005 (Abstract 2241)
17. Stenosis in the Coronary Artery of Normally Contracting Heart: Detection by a Simplified tissue Doppler Mapping Technique. *J Am Coll Cardiol* 2004; 53th: 371A (Abstract)