

- (1) Reduced Plasma Eicosapentaenoic Acid-Arachidonic Acid Ratio in Peripheral Artery Disease: *Angiology*. 2013 Feb; 64(2):112-8.
- (2) Short and mid-term results of balloon angioplasty for renal artery fibromuscular dysplasia: *Cardiovasc Interv Ther*. 2014 Oct; 29(4):293-9.
- (3) Iliac artery fibromuscular dysplasia successfully treated by balloon angioplasty guided by intravascular ultrasound and pressure wire measurements: A case report *Journal of cardiology cases*. 2014 May:203-207
- (4) Outcomes of Zilver PTX Stent Implantation for the Treatment of Complex Femoropopliteal Artery Disease: *Heart Vessels*. 2016 31(2):152-7.
- (5) Endovascular therapy by CO(2) angiography to prevent contrast-induced nephropathy in patients with chronic kidney disease: A prospective multicenter trial of CO(2) angiography registry. *Catheter Cardiovasc Interv*. 2015 85(5):870-7.
- (6) Clinical Outcomes of Renal Artery Stenting for Hypertension and Chronic Kidney Disease over 12 Months in J-RAS study: A Prospective, Single-Arm, Multi-center Clinical Study: *Circulation J* 2015;79(2):351-359
- (7) Nitinol stent implantation for femoropopliteal disease in patients on hemodialysis: results of the 3-year retrospective multicenter APOLLON study: *Heart and Vessels* 2015
- (8) Appropriate hemostasis by routine use of ultrasound echo-guided transfemoral access and vascular closure devices after lower extremity percutaneous revascularization. *Cardiovasc Interv Ther*. 2016 Jul 18.
- (9) Angiographic dissection patterns and patency outcomes of post balloon angioplasty for superficial femoral artery disease *J EVT* 2017